

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number  
**WO 03/013247 A1**

(51) International Patent Classification<sup>7</sup>: **A01N 43/40**,  
C07D 213/61, 213/64, 213/16, 213/73, 213/57, 213/75,  
213/65, 213/74

(21) International Application Number: PCT/EP02/08878

(22) International Filing Date: 8 August 2002 (08.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
1471/01 9 August 2001 (09.08.2001) CH

(71) Applicant (for all designated States except US): **SYNGENTA PARTICIPATIONS AG** [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SCHAETZER, Juergen** [DE/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **EBERLE, Martin** [CH/CH]; Asterhagstrasse 22, CH-4103 Bottmingen (CH). **WENGER, Jean** [CH/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **BERTEINA-RABOIN, Sabine** [FR/FR]; Université d'Orléans, UFR Sciences, I.C.O.A., Rue de Chartres, F-45067 Orléans (FR). **NEBEL, Kurt** [CH/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **STOLLER, André** [CH/FR]; 7, rue Charles Wolf, F-68730 Blotzheim (FR). **HALL, Roger, Graham** [GB/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **BONDY, Steven, Scott** [US/US]; Combichem, Deltagen Research Labs, Suite

400, 4570 Executive Drive, San Diego, CA 92121 (US). **COMER, Daniel, Dennis** [US/US]; Combichem, Deltagen Research Labs, Suite 400, 4570 Executive Drive, San Diego, CA 92121 (US). **PENZOTTI, Julie, Elizabeth** [US/US]; Combichem, Deltagen Research Labs, Suite 400, 4570 Executive Drive, San Diego, CA 92121 (US). **GROOTENHUIS, Peter, Diederik, Jan** [NL/US]; Combichem, Deltagen Research Labs, Suite 400, 4570 Executive Drive, San Diego, CA 92121 (US).

(74) Agent: **BASTIAN, Werner**; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRIDYLPROPYNYLOXYPHENYL DERIVATIVES FOR USE AS HERBICIDES

(57) Abstract: Compounds of formula (I), wherein the substituents R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and Z and the suffixes n and m are as defined in claim 1, and the agrochemically acceptable salts and all stereoisomers and tautomers of those compounds are suitable for use as herbicides.



WO 03/013247 A1

## PYRIDYLPROPYNLOXYPHENYL DERIVATIVES FOR USE AS HERBICIDES


Novel herbicides

The present invention relates to novel herbicidally active pyridyl-alkynes and pyridyl N-oxide-alkynes, to processes for their preparation, to compositions comprising those compounds, and to their use in controlling weeds, especially in crops of useful plants, or in inhibiting plant growth.

Phenylalkynes having herbicidal action are described, for example, in JP-A-11 147 866, WO 01/55066 and PCT Application No. EP01/11353.

Novel pyridyl-alkynes and pyridyl N-oxide-alkynes having herbicidal and growth-inhibiting properties have now been found.

The present invention accordingly relates to compounds of formula I



wherein

Z is =N- or  $\text{—}\overset{\text{||}}{\text{N}}^+\text{—O}^-$  ;

n is 0, 1, 2, 3, 4 or 5;

each  $R_1$  independently of any others is halogen, -CN, -SCN, -SF<sub>5</sub>, -NO<sub>2</sub>, -NR<sub>5</sub>R<sub>6</sub>, -CO<sub>2</sub>R<sub>7</sub>, -CONR<sub>8</sub>R<sub>9</sub>, -C(R<sub>10</sub>)=NOR<sub>11</sub>, -COR<sub>12</sub>, -OR<sub>13</sub>, -SR<sub>14</sub>, -SOR<sub>15</sub>, -SO<sub>2</sub>R<sub>16</sub>, -OSO<sub>2</sub>R<sub>17</sub>, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl; or is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl or C<sub>2</sub>-C<sub>8</sub>alkynyl substituted by one or more halogen, -CN, -NO<sub>2</sub>, -NR<sub>18</sub>R<sub>19</sub>, -CO<sub>2</sub>R<sub>20</sub>, -CONR<sub>21</sub>R<sub>22</sub>, -COR<sub>23</sub>, -C(R<sub>24</sub>)=NOR<sub>25</sub>, -C(S)NR<sub>26</sub>R<sub>27</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>28</sub>, -OR<sub>29</sub>, -SR<sub>30</sub>, -SOR<sub>31</sub>, -SO<sub>2</sub>R<sub>32</sub> or C<sub>3</sub>-C<sub>6</sub>cycloalkyl substituents; or

each  $R_1$  independently of any others is C<sub>3</sub>-C<sub>6</sub>cycloalkyl substituted by one or more halogen, -CN, -NO<sub>2</sub>, -NR<sub>18</sub>R<sub>19</sub>, -CO<sub>2</sub>R<sub>20</sub>, -CONR<sub>21</sub>R<sub>22</sub>, -COR<sub>23</sub>, -C(R<sub>24</sub>)=NOR<sub>25</sub>, -C(S)NR<sub>26</sub>R<sub>27</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>28</sub>, -SR<sub>30</sub>, -SOR<sub>31</sub>, -SO<sub>2</sub>R<sub>32</sub> or C<sub>3</sub>-C<sub>6</sub>cycloalkyl substituents; or

each  $R_1$  independently of any others is phenyl, which may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or

two adjacent  $R_1$  together form a  $C_1$ - $C_7$ alkylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9; or

two adjacent  $R_1$  together form a  $C_2$ - $C_7$ alkenylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9;

$R_3$  and  $R_4$  are each independently of the other hydrogen, halogen, -CN,  $C_1$ - $C_4$ alkyl or  $C_1$ - $C_4$ alkoxy; or

$R_3$  and  $R_4$  together are  $C_2$ - $C_5$ alkylene;

$R_5$  is hydrogen or  $C_1$ - $C_8$ alkyl;

$R_6$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl; wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or

$R_5$  and  $R_6$  together are a  $C_2$ - $C_5$ alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

$R_7$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl or  $C_3$ - $C_8$ alkynyl, or is  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl or  $C_3$ - $C_8$ alkynyl substituted by one or more halogen,  $C_1$ - $C_4$ alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents;

$R_8$  is hydrogen or  $C_1$ - $C_8$ alkyl;

$R_9$  is hydrogen or  $C_1$ - $C_8$ alkyl, or is  $C_1$ - $C_8$ alkyl substituted by one or more -COOH,  $C_1$ - $C_8$ alkoxycarbonyl or -CN substituents, or

$R_9$  is  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or

$R_8$  and  $R_9$  together are  $C_2$ - $C_5$ alkylene;

$R_{10}$  is hydrogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ cycloalkyl;

$R_{11}$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ haloalkenyl;

$R_{12}$  is hydrogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ cycloalkyl;

$R_{13}$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl or  $C_3$ - $C_8$ alkynyl; or

R<sub>13</sub> is phenyl or phenyl-C<sub>1</sub>-C<sub>6</sub>alkyl, wherein both phenyl rings may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl substituents, or

R<sub>13</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl substituted by one or more halogen, -CN, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)-amino or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>14</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>15</sub>, R<sub>16</sub> and R<sub>17</sub> are each independently of the others C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>18</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>19</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>18</sub> and R<sub>19</sub> together are a C<sub>2</sub>-C<sub>5</sub>alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

R<sub>20</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>21</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>22</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents, or

R<sub>22</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>21</sub> and R<sub>22</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>23</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>24</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>25</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>26</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>27</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents, or

R<sub>27</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>26</sub> and R<sub>27</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>28</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>29</sub> and R<sub>30</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>31</sub> and R<sub>32</sub> are each independently of the other C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

m is 0, 1, 2, 3 or 4;

each R<sub>2</sub> independently of any others is halogen, -CN, -SCN, -OCN, -N<sub>3</sub>, -SF<sub>5</sub>, -NO<sub>2</sub>, -NR<sub>33</sub>R<sub>34</sub>, -CO<sub>2</sub>R<sub>35</sub>, -CONR<sub>36</sub>R<sub>37</sub>, -C(R<sub>38</sub>)=NOR<sub>39</sub>, -COR<sub>40</sub>, -OR<sub>41</sub>, -SR<sub>42</sub>, -SOR<sub>43</sub>, -SO<sub>2</sub>R<sub>44</sub>, -OSO<sub>2</sub>R<sub>45</sub>, -N([CO]<sub>p</sub>R<sub>46</sub>)COR<sub>47</sub>, -N(OR<sub>54</sub>)COR<sub>55</sub>, -N(R<sub>56</sub>)SO<sub>2</sub>R<sub>57</sub>, -N(SO<sub>2</sub>R<sub>58</sub>)SO<sub>2</sub>R<sub>59</sub>, -N=C(OR<sub>60</sub>)R<sub>61</sub>, -CR<sub>62</sub>(OR<sub>63</sub>)OR<sub>64</sub>, -OC(O)NR<sub>65</sub>R<sub>66</sub>, -SC(O)NR<sub>67</sub>R<sub>68</sub>, -OC(S)NR<sub>69</sub>R<sub>70</sub> or -N-phthalimide; or

R<sub>2</sub> is a 5- to 7-membered heterocyclic ring system which may be aromatic or partially or fully saturated and may contain from 1 to 4 hetero atoms selected from nitrogen, oxygen and sulfur, it being possible for that heterocyclic ring system in turn to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, hydroxy-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>4</sub>alkyl, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl substituents;

R<sub>33</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl; and

R<sub>34</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>33</sub> and R<sub>34</sub> together are a C<sub>2</sub>-C<sub>5</sub>alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

R<sub>35</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>36</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>37</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents, or

R<sub>37</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>36</sub> and R<sub>37</sub> together are C<sub>3</sub>-C<sub>5</sub>alkylene;

R<sub>38</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>39</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>40</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>8</sub>alkylthio, -C(O)-C(O)OC<sub>1</sub>-C<sub>4</sub>alkyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl;

R<sub>41</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>8</sub>alkyl-carbonyl, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl, C<sub>3</sub>-C<sub>8</sub>alkenylloxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylthio-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl-C<sub>1</sub>-C<sub>6</sub>alkyl or C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl-C<sub>1</sub>-C<sub>6</sub>alkyl; or R<sub>41</sub> is phenyl or phenyl-C<sub>1</sub>-C<sub>6</sub>alkyl, wherein both phenyl rings may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, or -S(O)<sub>2</sub>C<sub>1</sub>-C<sub>8</sub>alkyl substituents, or

R<sub>41</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)amino or -CN substituents;

R<sub>42</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>43</sub> and R<sub>44</sub> are each independently of the other C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>45</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or

R<sub>45</sub> is phenyl, it being possible for the phenyl ring to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>8</sub>alkylthio, C<sub>1</sub>-C<sub>8</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>8</sub>alkylsulfonyl substituents;

R<sub>46</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or C<sub>1</sub>-C<sub>4</sub>haloalkyl;

R<sub>47</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, -NR<sub>48</sub>COR<sub>49</sub>, -NR<sub>50</sub>SO<sub>2</sub>R<sub>51</sub> or -NR<sub>52</sub>CO<sub>2</sub>R<sub>53</sub> substituents, or R<sub>47</sub> is phenyl or benzyl, each of which may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

p is 0 or 1;

R<sub>48</sub>, R<sub>49</sub>, R<sub>50</sub>, R<sub>51</sub>, R<sub>52</sub> and R<sub>53</sub> are each independently of the others hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, phenyl, benzyl or naphthyl, it being possible for the three last-mentioned aromatic radicals in turn to be substituted by one or more halogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino, -NH<sub>2</sub>, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>54</sub> and R<sub>55</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl or phenyl, whereby the phenyl ring may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>8</sub>alkylthio, C<sub>1</sub>-C<sub>8</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>8</sub>alkylsulfonyl substituents;

R<sub>56</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or benzyl, it being possible for benzyl in turn to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>8</sub>alkylthio, C<sub>1</sub>-C<sub>8</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>8</sub>alkylsulfonyl substituents;

R<sub>57</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, phenyl, benzyl or naphthyl, it being possible for the three last-mentioned aromatic rings to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino, -NH<sub>2</sub>, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>58</sub> and R<sub>59</sub> are each independently of the other C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl, benzyl or naphthyl, it being possible for the three last-mentioned aromatic rings to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino, -NH<sub>2</sub>, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>60</sub> and R<sub>61</sub> are each independently of the other hydrogen or C<sub>1</sub>-C<sub>6</sub>alkyl;

R<sub>62</sub>, R<sub>63</sub> and R<sub>64</sub> are each independently of the others hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or

R<sub>63</sub> and R<sub>64</sub> together form a C<sub>2</sub>-C<sub>5</sub>alkylene bridge;

R<sub>65</sub>, R<sub>66</sub>, R<sub>67</sub>, R<sub>68</sub>, R<sub>69</sub> and R<sub>70</sub> are each independently of the others hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or

R<sub>65</sub> and R<sub>66</sub> together or R<sub>67</sub> and R<sub>68</sub> together or R<sub>69</sub> and R<sub>70</sub> together form a C<sub>2</sub>-C<sub>5</sub>alkylene bridge; or

each R<sub>2</sub> independently of any others is C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl mono- or poly-substituted by halogen, -CN, -N<sub>3</sub>, -SCN, -NO<sub>2</sub>, -NR<sub>71</sub>R<sub>72</sub>, -CO<sub>2</sub>R<sub>73</sub>, -CONR<sub>74</sub>R<sub>75</sub>, -COR<sub>76</sub>, -C(R<sub>77</sub>)=NOR<sub>78</sub>, -C(S)NR<sub>79</sub>R<sub>80</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>81</sub>, -OR<sub>82</sub>, -SR<sub>83</sub>, -SOR<sub>84</sub>, -SO<sub>2</sub>R<sub>85</sub>, -O(SO<sub>2</sub>)R<sub>86</sub>, -N(R<sub>87</sub>)CO<sub>2</sub>R<sub>88</sub>, -N(R<sub>89</sub>)COR<sub>90</sub>, -S<sup>+</sup>(R<sub>91</sub>)<sub>2</sub>, -N<sup>+</sup>(R<sub>92</sub>)<sub>3</sub>, -Si(R<sub>93</sub>)<sub>3</sub> or C<sub>3</sub>-C<sub>6</sub>cycloalkyl; or

each R<sub>2</sub> independently of any others is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by a 5- to 7-membered heterocyclic ring system, which may be aromatic or partially or fully saturated and may contain from 1 to 4 hetero atoms selected from nitrogen, oxygen and sulfur, it being possible for that heterocyclic ring system in turn to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, hydroxy-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>4</sub>alkyl, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl substituents; or

each R<sub>2</sub> independently of any others is C<sub>2</sub>-C<sub>8</sub>alkenyl, or is C<sub>2</sub>-C<sub>8</sub>alkenyl mono- or poly-substituted by halogen, -CN, -NO<sub>2</sub>, -CO<sub>2</sub>R<sub>94</sub>, -CONR<sub>95</sub>R<sub>96</sub>, -COR<sub>97</sub>, -C(R<sub>98</sub>)=NOR<sub>99</sub>, -C(S)NR<sub>100</sub>R<sub>101</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>102</sub>, -OR<sub>103</sub>, -Si(R<sub>104</sub>)<sub>3</sub> or C<sub>3</sub>-C<sub>6</sub>cycloalkyl; or

each R<sub>2</sub> independently of any others is C<sub>2</sub>-C<sub>8</sub>alkynyl, or is C<sub>2</sub>-C<sub>8</sub>alkynyl mono- or poly-substituted by halogen, -CN, -CO<sub>2</sub>R<sub>105</sub>, -CONR<sub>106</sub>R<sub>107</sub>, -COR<sub>108</sub>, -C(R<sub>109</sub>)=NOR<sub>110</sub>, -C(S)NR<sub>111</sub>R<sub>112</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>113</sub>, -OR<sub>114</sub>, -Si(R<sub>115</sub>)<sub>3</sub> or C<sub>3</sub>-C<sub>6</sub>cycloalkyl; or

each  $R_2$  independently of any others is  $C_3$ - $C_6$ cycloalkyl, or is  $C_3$ - $C_6$ cycloalkyl mono- or poly-substituted by halogen, -CN,  $-CO_2R_{116}$ ,  $-CONR_{117}R_{118}$ ,  $-COR_{119}$ ,  $-C(R_{120})=NOR_{121}$ ,

$-C(S)NR_{122}R_{123}$  or  $-C(C_1-C_4\text{alkylthio})=NR_{124}$ ; or

two adjacent  $R_2$  together form a  $C_1$ - $C_7$ alkylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9; or

two adjacent  $R_2$  together form a  $C_2$ - $C_7$ alkenylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9;

$R_{71}$  is hydrogen or  $C_1$ - $C_8$ alkyl;

$R_{72}$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN,  $-NO_2$ ,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or

$R_{71}$  and  $R_{72}$  together are a  $C_2$ - $C_5$ alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

$R_{73}$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl or  $C_3$ - $C_8$ alkynyl, or is  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl or  $C_3$ - $C_8$ alkynyl substituted by one or more halogen,  $C_1$ - $C_4$ alkoxy or phenyl substituents, it being possible for phenyl in turn to be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN,  $-NO_2$ ,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents;

$R_{74}$  is hydrogen or  $C_1$ - $C_8$ alkyl;

$R_{75}$  is hydrogen,  $C_1$ - $C_8$ alkyl or  $C_3$ - $C_7$ cycloalkyl, or is  $C_1$ - $C_8$ alkyl substituted by one or more  $-COOH$ ,  $C_1$ - $C_8$ alkoxycarbonyl,  $C_1$ - $C_6$ alkoxy or  $-CN$  substituents; or

$R_{75}$  is  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN,  $-NO_2$ ,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or

$R_{74}$  and  $R_{75}$  together are a  $C_2$ - $C_5$ alkylene chain, which may be interrupted by an oxygen or sulfur atom;

$R_{76}$  is hydrogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ cycloalkyl;

$R_{77}$  is hydrogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ cycloalkyl;

$R_{78}$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ haloalkenyl; and

$R_{79}$  is hydrogen or  $C_1$ - $C_8$ alkyl;

$R_{80}$  is hydrogen or  $C_1$ - $C_8$ alkyl, or is  $C_1$ - $C_8$ alkyl substituted by one or more  $-COOH$ ,  $C_1$ - $C_8$ alkoxycarbonyl or  $-CN$  substituents; or



R<sub>80</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>79</sub> and R<sub>80</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>81</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>82</sub> is -Si(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>3</sub>, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or C<sub>1</sub>-C<sub>8</sub>alkyl, whereby C<sub>1</sub>-C<sub>8</sub>alkyl is mono- or poly-substituted by halogen, -CN, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)amino or C<sub>1</sub>-C<sub>4</sub>alkoxy;

R<sub>83</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or C<sub>1</sub>-C<sub>8</sub>alkyl, whereby C<sub>1</sub>-C<sub>8</sub>alkyl is mono- or poly-substituted by halogen, -CN, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)amino or C<sub>1</sub>-C<sub>4</sub>alkoxy;

R<sub>84</sub>, R<sub>85</sub> and R<sub>86</sub> are each independently of the others C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl which is substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>87</sub> and R<sub>89</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl or C<sub>1</sub>-C<sub>8</sub>alkoxy;

R<sub>88</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>90</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>91</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl;

R<sub>92</sub> and R<sub>93</sub> are each independently of the other C<sub>1</sub>-C<sub>6</sub>alkyl;

R<sub>94</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, each of which may be mono- or poly-substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>95</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>96</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents; or

R<sub>96</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>95</sub> and R<sub>96</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>97</sub> and R<sub>98</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>99</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>100</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>101</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents; or

R<sub>101</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>100</sub> and R<sub>101</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>102</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>103</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, -Si(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>3</sub>, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl;

R<sub>104</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl;

R<sub>105</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, each of which may be mono- or poly-substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>106</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>107</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl or -CN substituents; or

R<sub>107</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>106</sub> and R<sub>107</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>108</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>109</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>110</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>111</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>112</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl or -CN substituents; or

R<sub>112</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>111</sub> and R<sub>112</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>113</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>114</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, -Si(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>3</sub>, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl;

R<sub>115</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl;

R<sub>116</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, each of which may be mono- or poly-substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;


R<sub>117</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

$R_{118}$  is hydrogen or  $C_1$ - $C_8$ alkyl, or is  $C_1$ - $C_8$ alkyl substituted by one or more -COOH,  $C_1$ - $C_8$ alkoxycarbonyl or -CN substituents; or  
 $R_{118}$  is  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or  
 $R_{117}$  and  $R_{118}$  together are  $C_2$ - $C_5$ alkylene;  
 $R_{119}$  is hydrogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ cycloalkyl;  
 $R_{120}$  is hydrogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ cycloalkyl;  
 $R_{121}$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl,  $C_1$ - $C_4$ haloalkyl or  $C_3$ - $C_6$ haloalkenyl;  
 $R_{122}$  is hydrogen or  $C_1$ - $C_8$ alkyl;  
 $R_{123}$  is hydrogen or  $C_1$ - $C_8$ alkyl, or is  $C_1$ - $C_8$ alkyl substituted by one or more -COOH,  $C_1$ - $C_8$ alkoxycarbonyl or -CN substituents; or  
 $R_{123}$  is  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or  
 $R_{122}$  and  $R_{123}$  together are  $C_2$ - $C_5$ alkylene; and  
 $R_{124}$  is hydrogen or  $C_1$ - $C_8$ alkyl,  
and to the agrochemically acceptable salts and all stereoisomers and tautomers of the compounds of formula I.

When  $n$  is 0, all the free valencies on the phenyl ring of the compounds of formula I are substituted by hydrogen. When  $m$  is 0, all the free valencies on the pyridyl ring of the compounds of formula I are substituted by hydrogen.

Examples of substituents that are formed when  $R_5$  and  $R_6$  together or  $R_{18}$  and  $R_{19}$  together or  $R_{36}$  and  $R_{37}$  together or  $R_{74}$  and  $R_{75}$  together are a  $C_2$ - $C_5$ alkylene chain, which may be interrupted by an oxygen or a sulfur atom, are piperidine, morpholine, thiomorpholine and pyrrolidine.

Examples of heterocyclic ring systems, which may be aromatic or partially or fully saturated, in the definition of  $R_2$  are:



The alkyl groups appearing in the definitions of substituents may be straight-chain or branched and are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, and also the isomers of pentyl, hexyl, heptyl, octyl, nonyl and decyl.

Halogen is fluorine, chlorine, bromine and iodine, preferably fluorine and chlorine.

Haloalkyl is, for example, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 2-fluoroethyl, 2-chloroethyl, pentafluoroethyl, 1,1-difluoro-2,2,2-trichloroethyl, 2,2,3,3-tetrafluoroethyl and 2,2,2-trichloroethyl; preferably trichloromethyl, difluorochloromethyl, difluoromethyl, trifluoromethyl and dichlorofluoromethyl.

Alkoxy groups have preferably a chain length of from 1 to 6, especially from 1 to 4, carbon atoms. Alkoxy is, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy and tert-butoxy, and also the pentyloxy and hexyloxy isomers; preferably methoxy and ethoxy.




Alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, alkylsulfonyl, alkylsulfinyl, alkylaminoalkoxy, alkoxycarbonyl, alkylcarbonyloxy, alkenylthio, alkenylsulfonyl, alkenylsulfinyl, alkynylsulfonyl, alkynylthio and alkynylsulfinyl groups are derived from the mentioned alkyl radicals. The alkenyl and alkynyl groups can be mono- or poly-unsaturated. Alkenyl is to be understood as being, for example, vinyl, allyl, methallyl, 1-methylvinyl or but-2-en-1-yl. Alkynyl is, for example, ethynyl, propargyl, but-2-yn-1-yl, 2-methylbutyn-2-yl or but-3-yn-2-yl.

Alkylthio groups have preferably a chain length of from 1 to 4 carbon atoms. Alkylthio is, for example, methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, isobutylthio, sec-butylthio or tert-butylthio, preferably methylthio and ethylthio. Alkylsulfinyl is, for example, methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl, n-butylsulfinyl, isobutylsulfinyl, sec-butylsulfinyl or tert-butylsulfinyl; preferably methylsulfinyl or ethylsulfinyl.


Alkylsulfonyl is, for example, methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, isobutylsulfonyl, sec-butylsulfonyl or tert-butylsulfonyl; preferably methylsulfonyl or ethylsulfonyl.

Alkoxyalkyl groups have preferably from 1 to 6 carbon atoms. Alkoxyalkyl is, for example, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, n-propoxymethyl, n-propoxyethyl, isopropoxymethyl or isopropoxyethyl.

Substituents wherein two adjacent  $R_1$  together form a  $C_1$ - $C_7$ alkylene bridge which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9, or two adjacent  $R_1$  together form a  $C_2$ - $C_7$ alkenylene bridge which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9, have, for example, the following structures:



Substituents wherein two adjacent  $R_2$  together form a  $C_1$ - $C_7$ alkylene bridge which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9, or two adjacent  $R_2$  together form a  $C_2$ - $C_7$ alkenylene bridge which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9, have, for example, the following structures:



The invention relates also to the salts which the compounds of formula I are able to form especially with amines, alkali metal and alkaline earth metal bases or quaternary ammonium bases. Suitable salt-formers are described, for example, in WO 98/41089.

Among the alkali metal and alkaline earth metal hydroxides as salt formers, special mention should be made of the hydroxides of lithium, sodium, potassium, magnesium and calcium, but especially the hydroxides of sodium and potassium.

Examples of amines suitable for ammonium salt formation include ammonia as well as primary, secondary and tertiary C<sub>1</sub>-C<sub>18</sub>alkylamines, C<sub>1</sub>-C<sub>4</sub>hydroxyalkylamines and C<sub>2</sub>-C<sub>4</sub>-alkoxyalkylamines, for example methylamine, ethylamine, n-propylamine, isopropylamine, the four butylamine isomers, n-amylamine, isoamylamine, hexylamine, heptylamine, octylamine, nonylamine, decylamine, pentadecylamine, hexadecylamine, heptadecylamine, octadecylamine, methylethylamine, methylisopropylamine, methylhexylamine, methylnonylamine, methylpentadecylamine, methyloctadecylamine, ethylbutylamine, ethylheptylamine, ethyloctylamine, hexylheptylamine, hexyloctylamine, dimethylamine, diethylamine, di-n-propylamine, diisopropylamine, di-n-butylamine, di-n-amylamine, diisoamylamine, dihexylamine, diheptylamine, dioctylamine, ethanolamine, n-propanolamine, isopropanolamine, N,N-

diethanolamine, N-ethylpropanolamine, N-butylethanolamine, allylamine, n-butenyl-2-amine, n-pentenyl-2-amine, 2,3-dimethylbutenyl-2-amine, dibutenyl-2-amine, n-hexenyl-2-amine, propylenediamine, trimethylamine, triethylamine, tri-n-propylamine, triisopropylamine, tri-n-butylamine, triisobutylamine, tri-sec-butylamine, tri-n-amylamine, methoxyethylamine and ethoxyethylamine; heterocyclic amines, for example pyridine, quinoline, isoquinoline, morpholine, piperidine, pyrrolidine, indoline, quinuclidine and azepine; primary arylamines, for example anilines, methoxyanilines, ethoxyanilines, o-, m- and p-toluidines, phenylenediamines, benzidines, naphthylamines and o-, m- and p-chloroanilines; but especially triethylamine, isopropylamine and diisopropylamine.

Preferred quaternary ammonium bases suitable for salt formation correspond e.g. to the formula  $[N(R_a R_b R_c R_d)]OH$  wherein  $R_a$ ,  $R_b$ ,  $R_c$  and  $R_d$  are each independently of the other  $C_1$ - $C_4$ alkyl. Other suitable tetraalkylammonium bases with other anions can be obtained, for example, by anion exchange reactions.

Preferred compounds of formula I are those wherein Z is =N-; and each  $R_2$  independently of any others is  $C_2$ - $C_8$ alkenyl, or is  $C_2$ - $C_8$ alkenyl mono- or poly-substituted by -CN, -NO<sub>2</sub>, -CO<sub>2</sub>R<sub>94</sub>, -CONR<sub>95</sub>R<sub>96</sub>, -COR<sub>97</sub>, -C(R<sub>98</sub>)=NOR<sub>99</sub>, -C(S)NR<sub>100</sub>R<sub>101</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>102</sub>, -OR<sub>103</sub>, -Si(R<sub>104</sub>)<sub>3</sub> or  $C_3$ - $C_6$ cycloalkyl.


Further preferred compounds of formula I are those wherein each  $R_2$  independently of any others is halogen, -CN, -SCN, -OCN, -N<sub>3</sub>, -CONR<sub>36</sub>R<sub>37</sub>, -C(R<sub>38</sub>)=NOR<sub>39</sub>, -COR<sub>40</sub>, -OR<sub>41</sub>, -SO<sub>2</sub>R<sub>45</sub>, -N([CO]<sub>p</sub>R<sub>46</sub>)COR<sub>47</sub>, -N(R<sub>56</sub>)SO<sub>2</sub>R<sub>57</sub>, -N(SO<sub>2</sub>R<sub>58</sub>)SO<sub>2</sub>R<sub>59</sub>, -N=C(OR<sub>60</sub>)R<sub>61</sub> or  $C_1$ - $C_8$ alkyl, or is  $C_1$ - $C_8$ alkyl mono- or poly-substituted by halogen, -CN, -N<sub>3</sub>, -SCN, -CONR<sub>74</sub>R<sub>75</sub>, -COR<sub>76</sub>, -C(R<sub>77</sub>)=NOR<sub>78</sub>, -C(S)NR<sub>79</sub>R<sub>80</sub>, -OR<sub>82</sub>, -SOR<sub>84</sub>, -SO<sub>2</sub>R<sub>85</sub> or -N(R<sub>89</sub>)COR<sub>90</sub>.

Preference is likewise given to compounds of formula I wherein each  $R_1$  independently of any others is halogen, -CN,  $C_1$ - $C_3$ alkyl,  $C_1$ - $C_3$ haloalkyl,  $C_1$ - $C_3$ cyanoalkyl, -OR<sub>13</sub> or -C(R<sub>24</sub>)=NOR<sub>25</sub>;  $R_{13}$  is  $C_1$ - $C_3$ alkyl or di( $C_1$ - $C_4$ -alkyl)amino- $C_1$ - $C_4$ alkyl;  $R_{24}$  is hydrogen or methyl; and  $R_{25}$  is hydrogen or  $C_1$ - $C_3$ alkyl.


Also of importance are compounds of formula I wherein  $R_3$  and  $R_4$  are each independently of the other hydrogen or methyl.




The compounds of formula I can be prepared by methods known *per se* described, for example, in Tetrahedron 1997 (53), 12621-12628; Helv. Chim. Acta 2000 (83), 650-657; J. Chem. Res., Synop. 1996 (10), 462-463; Org. Prep. Proc. Int. 1995 (27), 129-160; Tetrahedron Organic Chemistry 2000 (20), 209-213; and K. Sonogashira in "Comprehensive Organic Synthesis", Editors I. Fleming *et al.*, Pergamon, Oxford 1991, Vol. 3, page 521 ff., for example by reacting a compound of formula II




wherein  $R_1$  and  $n$  are as defined for formula I, in the presence of a base, with a compound of formula III



wherein  $R_3$  and  $R_4$  are as defined for formula I and  $X_1$  is O-tosyl, O-mesyl, chlorine, bromine or iodine, to form a compound of formula IV



wherein  $R_1$ ,  $R_3$ ,  $R_4$  and  $n$  are as defined, and then coupling that compound with a compound of formula V or Va




wherein  $R_2$  and  $m$  are as defined for formula I and  $A$  is a leaving group, e.g. halogen or trifluoromethanesulfonate, in the presence of a palladium catalyst, and, if desired, oxidising

the resulting pyridine derivative of formula I wherein Z is =N- to form the corresponding

pyridine N-oxide of formula I wherein Z is  $\text{—}\overset{\overset{\text{||}}{\text{+}}}{\text{N}}\text{—}\overset{\text{—}}{\text{O}}\text{—}$ .

The preparation of the compounds of formula I can be carried out e.g. according to the individual Schemes 1, 2, 3, 4 and 5. For the individual synthesis schemes it is generally true that various substituents  $\text{R}_2$  in a compound of formula V or Va are either already present at the outset or can be introduced in succession, for example by nucleophilic or electrophilic aromatic substitution.

Similarly, the compound of formula V may at the outset already be in the form of the pyridine N-oxide derivative of formula Va



If desired, however, the N-oxide function can be introduced into the pyridyl ring of the compound of formula I wherein Z is =N- only at the end of the synthesis sequence, *via* oxidation by conventional methods, e.g. with hydrogen peroxide or organic peracids.

According to Reaction Scheme 1, the compounds of formula I can be obtained, for example, from substituted phenyl propargyl ethers of formula IV.


The propargyl ethers of formula IV can be obtained beforehand by etherification of phenols of formula II, which are reacted in the presence of a base with acetylene derivatives of formula III. Such etherification reactions are standard procedures and can be carried out e.g. analogously to Tetrahedron 1997 (53), 12621-12628; Helv. Chim. Acta 2000 (83), 650-657; and J. Chem. Res., Synop. 1996 (10), 462-463.

In the next step, the propargyl ethers of formula IV are coupled with substituted pyridine or pyridine N-oxide derivatives of formula V or Va, respectively, under typical Sonogashira conditions (K. Sonogashira in "Comprehensive Organic Synthesis", Editors I. Fleming *et al.*, Pergamon, Oxford 1991, Vol. 3, page 521 ff.; J. Org. Chem. 1998 (63), 8551-8553). Catalyst mixtures that come into consideration are, for example, tetrakis(triphenylphosphine)-palladium or bis(triphenylphosphine)-palladium dichloride together with copper iodide, and bases that


come into consideration (for the reductive elimination) are especially amines, for example triethylamine, diethylamine and diisopropylethylamine.

The pyridines or pyridine N-oxides of formula V or Va, respectively, preferably carry a leaving group A, wherein A is e.g. halogen or trifluoromethanesulfonate (Tetrahedron Organic Chemistry 2000 (20), 209-213; J. Org. Chem. 1997 (62), 1491-1500). As solvents for the Sonogashira reaction there are customarily used ethers, for example tetrahydrofuran, chlorinated hydrocarbons, for example chloroform, or dipolar aprotic solvents, for example dimethylformamide or dimethyl sulfoxide, or amines, for example triethylamine or piperidine.


Scheme 1



The Pd-catalysed cross-coupling of suitably substituted pyridine or pyridine N-oxide derivatives of formula V or Va, respectively, with propargyl alcohols or terminal acetylenes of formula VI



wherein  $R_3$  and  $R_4$  are as defined for formula I, is known generally as the Sonogashira reaction and is shown diagrammatically in Reaction Scheme 2 for the pyridine derivatives of formula V. That reaction is documented in detail in *Tetrahedron Organic Chemistry* 2000 (20), 209-213 and can be used for the preparation of the pyridyl and pyridyl N-oxide propargyl alcohols of formula VII




wherein  $R_2$ ,  $R_3$ ,  $R_4$ ,  $Z$  and  $m$  are as defined for formula I.

The activation of the alcohol of formula VII ( $Z$  is =N-) is carried out e.g. by sulfonylation or halogenation according to Scheme 2. The sulfonylation of the alcohol of formula VII is a standard reaction and can be carried out e.g. with a sulfonic acid chloride, for example mesyl chloride (MsCl) or para-toluenesulfonic acid chloride (p-TsCl), in the presence of a tertiary amine, for example triethylamine, or an aromatic amine, for example pyridine, in a solvent, e.g. a chlorinated hydrocarbon, for example carbon tetrachloride or methylene chloride, or an amine, for example pyridine. Such reactions are generally known and are described e.g. in *J. Org. Chem.* 1997 (62), 8987; *J. Het. Chem.* 1995 (32), 875-882; and also in *Tetrahedron Lett.* 1997 (38), 8671-8674.

The halogenation of the alcohol of formula VII ( $Z$  is =N-) can be carried out analogously to standard procedures. For example, the bromination is carried out with carbon tetrabromide in the presence of triphenylphosphine (*Synthesis* 1998, 1015-1018) in methylene chloride. The chlorination is carried out with mineral acids, for example with concentrated hydrochloric acid (*J. Org. Chem.* 1955 (20), 95) or with para-toluenesulfonic acid chloride in the presence of an amine, for example triethylamine in a solvent, e.g. methylene chloride (*Tetrahedron Lett.* 1984 (25), 2295).


The preparation of the pyridyl-propynyloxy-benzenes of formula I ( $Z$  is =N-) can be carried out analogously to *Synthesis* 1995, 707-712; and *Tetrahedron Lett.* 1994 (35), 6405-6408 by means of copper-iodide-catalysed etherification of the phenol of formula II in the presence of the tosylate or mesylate or halide of formula VIII (according to Scheme 2). Suitable solvents are dimethylformamide and acetonitrile, and suitable bases are especially potassium carbonate and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU).

Scheme 2




Compounds of formula I can also be obtained by further methods (according to Scheme 3).


Scheme 3




Accordingly, acetylene esters of formula X




wherein R<sub>2</sub>, Z and m are as defined for formula I, can be obtained, by means of Sonogashira coupling, from the compounds of formula IX



and activated pyridine derivatives of formula V or Va



wherein  $R_2$  and  $m$  are as defined and  $A$  is a leaving group as described above, analogously to Synthetic Communic. 1998 (28), 327-335. The esters of formula X can then be reduced or reacted with organometallic compounds, for example Grignard reagents, to form the alcohols of formula VII




wherein  $R_2$ ,  $Z$  and  $m$  are as defined for formula I and  $R_3$  and  $R_4$  are each independently of the other hydrogen,  $C_1$ - $C_4$ alkyl or  $C_1$ - $C_4$ alkoxy.

The reduction of the acetylene esters of formula X ( $Z$  is =N-) to the alcohols of formula VII ( $Z$  is =N-) can be carried out especially with hydrides by standard methods, for example with lithium aluminium hydride or sodium borohydride in a solvent, e.g. an ether, for example diethyl ether, dioxane or tetrahydrofuran, or an alcohol, for example methanol or ethanol. Such reductions are described e.g. in C. Ferri, "Reaktionen der organischen Synthese" 1978, pages 98-102.


Reactions of carboxylic acid esters with Grignard reagents are standard in organic synthesis chemistry and are described in detail in "Organikum" 1976, pages 617-625. The subsequent etherification of the phenol derivatives of formula II in the presence of a compound of formula VIII to form the compounds of formula I has already been described in detail in Scheme 2.

Further methods of preparing the desired compounds of formula I are shown in Scheme 4 (variant of Scheme 3).


Scheme 4



Accordingly, a pyridylacetylene of formula XI



wherein  $\text{R}_2$  and  $m$  are as defined for formula I, is reacted with *n*-butyllithium (*n*-BuLi) and then with a chloroformic acid methyl ester to form an ester of formula Xa




wherein  $\text{Z}$  is =N-.


That ester can be converted into the desired compound of formula I entirely analogously to the method already described in Scheme 3, *via* an alcohol of formula VII ( $\text{Z}$  is =N-) (analogously to J. Org. Chem. 1988 (53), 4166-4171).

The compounds of formula I can also be prepared by first reacting the propargyl alcohols of formula VI







wherein  $\text{R}_3$  and  $\text{R}_4$  are as defined for formula I, with activated phenyl halides of formula XII



wherein  $\text{X}_2$  is halogen,  $n$  is 1, 2, 3, 4 or 5 and  $\text{R}_1$  is a substituent having an electron-withdrawing effect ( $-\text{M}$  and/or  $-\text{I}$  effect), e.g.  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $\text{CF}_3$  or  $\text{COR}_{12}$ , to form compounds of formula IV




wherein  $\text{R}_1$ ,  $\text{R}_3$ ,  $\text{R}_4$  and  $n$  are as defined, and then in the next synthesis step carrying out a Sonogashira reaction with activated pyridine or pyridine N-oxide derivatives of formula V or Va



wherein  $\text{R}_2$  and  $m$  are as defined for formula I and  $\text{A}$  is a leaving group, e.g. halogen or trifluoromethanesulfonate (Reaction Scheme 5).

## Scheme 5



The following comments apply to the individual reaction steps in Schemes 1 to 5:

The reactions to form compounds of formula I are advantageously performed in aprotic, inert organic solvents. Such solvents are hydrocarbons, such as benzene, toluene, xylene or cyclohexane, chlorinated hydrocarbons, such as dichloromethane, trichloromethane, tetrachloromethane and chlorobenzene, ethers, such as diethyl ether, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether, tetrahydrofuran and dioxane, nitriles, such as acetonitrile and propionitrile, amides, such as N,N-dimethylformamide, diethylformamide and N-methylpyrrolidinone. The reaction temperatures are preferably from -20°C to +120°C. The reactions generally proceed slightly exothermically and can generally be carried out at room temperature. In order to shorten the reaction time or alternatively to initiate the reaction, the reaction mixture may, if appropriate, be heated to its boiling point for a short time. The reaction times may likewise be shortened by the addition of a few drops of base as reaction catalyst. Suitable bases are especially tertiary amines, such as trimethylamine, triethylamine, quinuclidine, 1,4-diazabicyclo[2.2.2]octane, 1,5-diazabicyclo[4.3.0]non-5-ene and 1,5-diazabicyclo[5.4.0]undec-7-ene, but it is also possible to use inorganic bases, such as hydrides,

e.g. sodium or calcium hydride, hydroxides, such as sodium or potassium hydroxide, carbonates, such as sodium or potassium carbonate, or hydrogen carbonates, such as potassium or sodium hydrogen carbonate.

The compounds of formula I can be isolated in customary manner by concentration and/or evaporation of the solvent and can be purified by recrystallisation or trituration of the solid residue in solvents in which they are not readily soluble, such as ethers, aromatic hydrocarbons or chlorinated hydrocarbons.

The starting compounds of formulae II, III, V, VI, IX, XI and XII used in Schemes 1 to 5 are known, in some cases are commercially available or can be prepared analogously to described standard methods. For example, the compounds of formula V are described in Tetrahedron Organic Chemistry 20, 209 (2000).

For the use according to the invention of the compounds of formula I, or of compositions comprising them, there come into consideration all methods of application customary in agriculture, for example pre-emergence application, post-emergence application and seed dressing, and also various methods and techniques such as, for example, the controlled release of active ingredient. For that purpose a solution of the active ingredient is applied to mineral granule carriers or polymerised granules (urea/formaldehyde) and dried. If required, it is also possible to apply a coating (coated granules), which allows the active ingredient to be released in metered amounts over a specific period of time.

The compounds of formula I may be used as herbicides in their unmodified form, that is to say as obtained in the synthesis, but they are preferably formulated in customary manner together with the adjuvants conventionally employed in formulation technology, for example into emulsifiable concentrates, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules or microcapsules. Such formulations are described, for example, on pages 9 to 13 of WO 97/34485. As with the nature of the compositions, the methods of application, such as spraying, atomising, dusting, wetting, scattering or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances.

The formulations, that is to say the compositions, preparations or mixtures comprising the compound (active ingredient) of formula I or at least one compound of formula I and, usually, one or more solid or liquid formulation adjuvants, are prepared in known manner, e.g. by

homogeneously mixing and/or grinding the active ingredients with the formulation adjuvants, for example solvents or solid carriers. Surface-active compounds (surfactants) may also be used in addition in the preparation of the formulations. Examples of solvents and solid carriers are given, for example, on page 6 of WO 97/34485.

Depending upon the nature of the compound of formula I to be formulated, suitable surface-active compounds are non-ionic, cationic and/or anionic surfactants and surfactant mixtures having good emulsifying, dispersing and wetting properties. Examples of suitable anionic, non-ionic and cationic surfactants are listed, for example, on pages 7 and 8 of WO 97/34485. In addition, the surfactants conventionally employed in formulation technology, which are described, *inter alia*, in "McCutcheon's Detergents and Emulsifiers Annual" MC Publishing Corp., Ridgewood New Jersey, 1981, Stache, H., "Tensid-Taschenbuch", Carl Hanser Verlag, Munich/Vienna 1981, and M. and J. Ash, "Encyclopedia of Surfactants", Vol. I-III, Chemical Publishing Co., New York, 1980-81, are also suitable for the preparation of the herbicidal compositions according to the invention.

The herbicidal formulations generally contain from 0.1 to 99 % by weight, especially from 0.1 to 95 % by weight, of herbicide, from 1 to 99.9 % by weight, especially from 5 to 99.8 % by weight, of a solid or liquid formulation adjuvant, and from 0 to 25 % by weight, especially from 0.1 to 25 % by weight, of a surfactant. Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations. The compositions may also comprise further ingredients, such as stabilisers, for example vegetable oils or epoxidised vegetable oils (epoxidised coconut oil, rapeseed oil or soybean oil), anti-foams, for example silicone oil, preservatives, viscosity regulators, binders, tackifiers, and also fertilisers or other active ingredients.

The compounds of formula I are generally applied to plants or the locus thereof at rates of application of from 0.001 to 4 kg/ha, especially from 0.005 to 2 kg/ha. The concentration required to achieve the desired effect can be determined by experiment. It is dependent on the nature of the action, the stage of development of the cultivated plant and of the weed and on the application (place, time, method) and may vary within wide limits as a function of those parameters.


The compounds of formula I are distinguished by herbicidal and growth-inhibiting properties, allowing them to be used in crops of useful plants, especially cereals, cotton, soybeans, sugar beet, sugar cane, plantation crops, rape, maize and rice, and also for non-selective

weed control. The term "crops" is to be understood as including also crops that have been made tolerant to herbicides or classes of herbicides as a result of conventional methods of breeding or genetic techniques. The weeds to be controlled may be either monocotyledonous or dicotyledonous weeds, such as, for example, *Stellaria*, *Nasturtium*, *Agrostis*, *Digitaria*, *Avena*, *Setaria*, *Sinapis*, *Lolium*, *Solanum*, *Echinochloa*, *Scirpus*, *Monochoria*, *Sagittaria*, *Bromus*, *Alopecurus*, *Sorghum halepense*, *Panicum*, *Rottboellia*, *Cyperus*, *Abutilon*, *Sida*, *Xanthium*, *Amaranthus*, *Chenopodium*, *Ipomoea*, *Chrysanthemum*, *Galium*, *Viola* and *Veronica*.

The following Examples further illustrate but do not limit the invention.


Preparation Examples:

Example P1: Preparation of 3-methoxy-4-prop-2-ynyloxy-benzaldehyde O-methyl-oxime




5.0 g (26.3 mmol) of 3-methoxy-4-(2-propynyloxy)-benzaldehyde (see DE-A-4 141 401) are dissolved at 20°C in 20 ml of ethanol under nitrogen. Then, with stirring, 2.86 g (34.3 mmol) of O-methyl-hydroxylamine hydrochloride and 4.65 g (34.2 mmol) of anhydrous sodium acetate are added in succession thereto. After the addition, stirring is carried out for a further 18 hours at 20°C and 1.5 hours at about 50°C. The solvent is then distilled off, 100 ml of water are added to the residue and extraction is carried out three times with a total of 100 ml of dichloromethane. The combined organic phases are dried over magnesium sulfate. After evaporating off the solvent, 5.37 g of the desired target compound 3-methoxy-4-prop-2-ynyloxy-benzaldehyde O-methyl-oxime are obtained in the form of yellow crystals having a melting point of 68-69°C.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ (ppm) = 2.53 (t); 3.92 (s); 3.97 (s); 4.80 (t); 7.00 (s); 7.29 (s); 8.00 (s).

Example P2: Preparation of 4-fluoro-2-methoxy-1-prop-2-ynyloxy-benzene


80.0 g (0.563 mol) of 4-fluoro-2-methoxyphenol are dissolved at 20°C in 2 litres of acetone. 80.0 g of potassium carbonate are added and stirring is carried out at 20°C for 1 hour. Then, in the course of 30 minutes, 82.7 ml of propargyl bromide are added dropwise, with stirring, and the resulting suspension is heated at reflux temperature. When the reaction is complete, the solvent is distilled off and the residue is taken up in ether. The ether phase is washed three times with 1N NaOH, twice with water and twice with saturated brine. A small amount of toluene is then added to the ether phase and the reaction mixture is finally completely concentrated by evaporation. 171.6 g of the desired target compound 4-fluoro-2-methoxy-1-prop-2-ynyloxy-benzene are obtained in the form of a light-brown oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ (ppm) = 2.52 (s); 3.86 (s); 4.72 (s); 6.58-6.72 (m); 6.95-7.05 (m).


Example P3: 2-Chloro-5-iodopyridine

22.1 g (0.1 mol) of 2-hydroxy-5-iodo-pyridine are heated together with 31.0 g (0.2 mol) of phosphorus oxytrichloride (POCl<sub>3</sub>) for 1 hour at reflux temperature. When the reaction is complete, excess POCl<sub>3</sub> is distilled off and the residue is taken up in toluene. The organic phase is stirred with aqueous potassium carbonate solution, separated and concentrated by evaporation. The crude product is purified by chromatography over silica gel. 19 g of the desired title compound are obtained in the form of colourless crystals.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ (ppm) = 7.10-7.20 (d); 7.90-8.00 (dxd); 8.55-8.65 (d).

Example P4: 2-Chloro-5-[3-(4-fluoro-2-methoxy-phenoxy)-prop-1-ynyl]-pyridine

300 mg (1.25 mmol) of 2-chloro-5-iodo-pyridine (Example P3), 339 mg (1.87 mmol) of 4-fluoro-2-methoxy-1-prop-2-ynyloxy-benzene (Example P2) and 48 mg (0.25 mmol) of copper(I) iodide (CuI) are suspended in a mixture consisting of 4 ml of dioxane and 3 ml of diisopropylamine under argon at 20°C. The resulting reaction mixture is heated to 50°C and 88 mg (0.125 mmol) of Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> are added. After 3.5 hours, the reaction mixture is cooled to 20°C. The solvent mixture is distilled off *in vacuo* and the crude product is subjected to flash chromatography over silica gel (eluant: ethyl acetate/petroleum ether 1/5). 308 mg of the desired target compound 2-chloro-5-[3-(4-fluoro-2-methoxy-phenoxy)-prop-1-ynyl]-pyridine are obtained in the form of a beige solid having a melting point of 86-87°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ (ppm) = 3.87 (s); 4.93 (s); 6.56-6.70 (m); 6.97-7.02 (dxd); 7.28 (d); 7.64 (dxd); 8.42 (d).


Example P5: 2-[3-(4-Fluoro-2-methoxy-phenoxy)-prop-1-ynyl]-5-methyl-pyridine

200 mg (1.16 mmol) of 2-bromo-5-methyl-pyridine, 314 mg (1.74 mmol) of 4-fluoro-2-methoxy-1-prop-2-ynyloxy-benzene (Example P2) and 44 mg (0.23 mmol) of copper(I) iodide (CuI) are suspended in a mixture consisting of 4 ml of dioxane and 3 ml of diisopropylamine under argon at 20°C. The reaction mixture is heated to 50°C and 81 mg (0.12 mmol) of Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> are added. After 4 hours, the reaction mixture is cooled to 20°C. The solvent mixture is distilled off *in vacuo* and the resulting crude product is purified by chromatography over silica gel (eluant: ethyl acetate/petroleum ether 1/3). 208 mg of the desired target com-

pound 2-[3-(4-fluoro-2-methoxy-phenoxy)-prop-1-ynyl]-5-methyl-pyridine are obtained in the form of a brown oil.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  (ppm) = 2.33 (s); 3.86 (s); 4.95 (s); 6.55-6.68 (m); 7.05 (dxd); 7.29 (d); 7.43 (dxd); 8.40 (d).

Example P6: 2-[3-(4-Fluoro-2-methoxy-phenoxy)-prop-1-ynyl]-4-methyl-pyridine




200 mg (1.16 mmol) of 2-bromo-4-methyl-pyridine, 314 mg (1.74 mmol) of 4-fluoro-2-methoxy-1-prop-2-ynyloxy-benzene (Example P2) and 44 mg (0.23 mmol) of copper(I) iodide (CuI) are suspended in a mixture consisting of 4 ml of dioxane and 3 ml of diisopropylamine under argon at 20°C. The reaction mixture is heated to 50°C and 81 mg (0.12 mmol) of  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$  are added. After 4 hours, the reaction mixture is cooled to 20°C. The solvent mixture is distilled off *in vacuo* and the resulting crude product is purified by chromatography over silica gel (eluant: ethyl acetate/petroleum ether 1/3). 152 mg of the desired target compound 2-[3-(4-fluoro-2-methoxy-phenoxy)-prop-1-ynyl]-4-methyl-pyridine are obtained in the form of a brown solid.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  (ppm) = 2.32 (s); 3.87 (s); 4.95 (s); 6.56-6.68 (m); 7.03-7.08 (m); 7.23 (s); 8.41 (d).

In a manner analogous to that described in Examples P1 to P5 or in accordance with the methods as shown in Reaction Schemes 1-5 and in the references indicated, it is also possible to obtain the preferred compounds listed in the following Tables. In the column headed "Phys. data", the temperatures indicate the melting point (m.p.) of the compounds in question. In cases where the purity of the compounds has been investigated by means of HPLC/MS ("High Pressure Liquid Chromatography/Electrospray Mass Spectrometry"), the column headed "Phys. data" gives the  $[\text{M}+\text{H}]^+$  peak from the Electrospray-MS of the compound in question (e.g. Comp. No. 3.011).



Table 1: Compounds of formula I<sub>1</sub>


Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
1.001	2-OCH <sub>3</sub> , 4-CN	2-Cl	H	H	160-161
1.002	2-F, 4-Cl	2-Cl	H	H	
1.003	2-Cl, 4-Cl	2-Cl	H	H	
1.004	2-OCH <sub>3</sub> , 4-F	2-Cl	H	H	86-87
1.005	2-OCH <sub>3</sub> , 4-Cl	2-Cl	H	H	
1.006	2-OCH <sub>3</sub> , 4-Br	2-Cl	H	H	
1.007	2-CF <sub>3</sub> , 4-F	2-Cl	H	H	
1.008	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-Cl	H	H	
1.009	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-Cl	H	H	
1.010	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-Cl	H	H	97-99
1.011	2-OCH <sub>3</sub> , 5-CH=NOCH <sub>3</sub>	2-Cl	H	H	128-129
1.012	3-CF <sub>3</sub>	2-OCH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	H	H	oil
1.013	4-OCH <sub>3</sub>	2-OCH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	H	H	oil
1.014	H	2-OCH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	H	H	oil
1.015	2-Cl	2-OCH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	H	H	oil
1.016	4-Cl	2-OCH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	H	H	oil
1.017	3-Cl	2-OCH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	H	H	oil
1.018	2-OCH <sub>3</sub> , 4-F	H	H	H	78-79
1.019	2-OCH <sub>3</sub> , 4-CN	2-Cl	CH <sub>3</sub>	H	-
1.020	2-F, 4-Cl	2-Cl	CH <sub>3</sub>	H	-
1.021	2-Cl, 4-Cl	2-Cl	CH <sub>3</sub>	H	-
1.022	2-OCH <sub>3</sub> , 4-F	2-Cl	CH <sub>3</sub>	H	-
1.023	2-OCH <sub>3</sub> , 4-Cl	2-Cl	CH <sub>3</sub>	H	-
1.024	2-OCH <sub>3</sub> , 4-Br	2-Cl	CH <sub>3</sub>	H	-
1.025	2-CF <sub>3</sub> , 4-F	2-Cl	CH <sub>3</sub>	H	-
1.026	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-Cl	CH <sub>3</sub>	H	-
1.027	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-Cl	CH <sub>3</sub>	H	-

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
1.028	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-Cl	CH <sub>3</sub>	H	-
1.029	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-NH <sub>2</sub>	H	H	135-138
1.030	2-OCH <sub>3</sub> , 4-F	2-NH <sub>2</sub>	H	H	-
1.031	2-OCH <sub>3</sub> , 4-Cl	2-NH <sub>2</sub>	H	H	-
1.032	2-OCH <sub>3</sub> , 4-CN	3-Br	H	H	-
1.033	2-F, 4-Cl	3-Br	H	H	-
1.034	2-Cl, 4-Cl	3-Br	H	H	-
1.035	2-OCH <sub>3</sub> , 4-F	3-Br	H	H	72-74
1.036	2-OCH <sub>3</sub> , 4-Cl	3-Br	H	H	-
1.037	2-OCH <sub>3</sub> , 4-Br	3-Br	H	H	-
1.038	2-CF <sub>3</sub> , 4-F	3-Br	H	H	-
1.039	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	3-Br	H	H	-
1.040	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	3-Br	H	H	-
1.041	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-Br	H	H	102-104
1.042	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-Br, 6-OH	H	H	crystalline
1.043	2-OCH <sub>3</sub> , 4-F	3-Br, 6-OH	H	H	crystalline
1.044	2-OCH <sub>3</sub> , 4-CN	3-CH <sub>2</sub> CN	H	H	-
1.045	2-F, 4-Cl	3-CH <sub>2</sub> CN	H	H	-
1.046	2-Cl, 4-Cl	3-CH <sub>2</sub> CN	H	H	-
1.047	2-OCH <sub>3</sub> , 4-F	3-CH <sub>2</sub> CN	H	H	-
1.048	2-OCH <sub>3</sub> , 4-Cl	3-CH <sub>2</sub> CN	H	H	-
1.049	2-OCH <sub>3</sub> , 4-Br	3-CH <sub>2</sub> CN	H	H	-
1.050	2-CF <sub>3</sub> , 4-F	3-CH <sub>2</sub> CN	H	H	-
1.051	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	3-CH <sub>2</sub> CN	H	H	-
1.052	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	3-CH <sub>2</sub> CN	H	H	-
1.053	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-CH <sub>2</sub> CN	H	H	-
1.054	2-OCH <sub>3</sub> , 4-F	3-OCH <sub>3</sub> , 6-NHC(O)O-t-C <sub>4</sub> H <sub>9</sub>	H	H	crystalline
1.055	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-OCH <sub>3</sub> , 6-NHC(O)O-t-C <sub>4</sub> H <sub>9</sub>	H	H	crystalline
1.056	2-OCH <sub>3</sub> , 4-F	3-OCH <sub>3</sub> , 6-NH <sub>2</sub>	H	H	amorphous
1.057	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-OCH <sub>3</sub> , 6-NH <sub>2</sub>	H	H	crystalline
1.058	2-OCH <sub>3</sub> , 4-CN	3-Cl	H	H	-
1.059	2-F, 4-Cl	3-Cl	H	H	-

- 34 -

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
1.060	2-Cl, 4-Cl	3-Cl	H	H	-
1.061	2-OCH <sub>3</sub> , 4-F	3-Cl	H	H	-
1.062	2-OCH <sub>3</sub> , 4-Cl	3-Cl	H	H	-
1.063	2-OCH <sub>3</sub> , 4-Br	3-Cl	H	H	-
1.064	2-CF <sub>3</sub> , 4-F	3-Cl	H	H	-
1.065	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	3-Cl	H	H	-
1.066	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	3-Cl	H	H	-
1.067	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-Cl	H	H	-
1.068	2-OCH <sub>3</sub> , 4-F	3-Cl, 6-OH	H	H	-
1.069	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-Cl, 6-OH	H	H	crystalline
1.070	2-OCH <sub>3</sub> , 4-CN	3-CH(CH <sub>3</sub> )CN	H	H	-
1.071	2-F, 4-Cl	3-CH(CH <sub>3</sub> )CN	H	H	-
1.072	2-Cl, 4-Cl	3-CH(CH <sub>3</sub> )CN	H	H	-
1.073	2-OCH <sub>3</sub> , 4-F	3-CH(CH <sub>3</sub> )CN	H	H	-
1.074	2-OCH <sub>3</sub> , 4-Cl	3-CH(CH <sub>3</sub> )CN	H	H	-
1.075	2-OCH <sub>3</sub> , 4-Br	3-CH(CH <sub>3</sub> )CN	H	H	-
1.076	2-CF <sub>3</sub> , 4-F	3-CH(CH <sub>3</sub> )CN	H	H	-
1.077	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	3-CH(CH <sub>3</sub> )CN	H	H	-
1.078	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	3-CH(CH <sub>3</sub> )CN	H	H	-
1.079	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-CH(CH <sub>3</sub> )CN	H	H	-
1.080	2-OCH <sub>3</sub> , 4-F	3-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
1.081	2-OCH <sub>3</sub> , 4-Cl	3-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
1.082	2-OCH <sub>3</sub> , 4-Br	3-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
1.083	2-OCH <sub>3</sub> , 4-CN	3-CH <sub>3</sub>	H	H	-
1.084	2-F, 4-Cl	3-CH <sub>3</sub>	H	H	-
1.085	2-Cl, 4-Cl	3-CH <sub>3</sub>	H	H	-
1.086	2-OCH <sub>3</sub> , 4-F	3-CH <sub>3</sub>	H	H	-
1.087	2-OCH <sub>3</sub> , 4-Cl	3-CH <sub>3</sub>	H	H	-
1.088	2-OCH <sub>3</sub> , 4-Br	3-CH <sub>3</sub>	H	H	-
1.089	2-CF <sub>3</sub> , 4-F	3-CH <sub>3</sub>	H	H	-
1.090	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	3-CH <sub>3</sub>	H	H	-
1.091	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	3-CH <sub>3</sub>	H	H	-
1.092	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-CH <sub>3</sub>	H	H	-
1.093	2-OCH <sub>3</sub>	3-CH <sub>2</sub> CN	H	H	-

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
1.094	2-OCH <sub>3</sub>	4-CH <sub>2</sub> CN	H	H	-
1.095	2-OCH <sub>3</sub>	3-F	H	H	-
1.096	2-OCH <sub>3</sub>	3-Cl	H	H	-
1.097	2-OCH <sub>3</sub>	3-Br	H	H	-
1.098	2-OCH <sub>3</sub> , 4-F	2-OCH <sub>3</sub>	H	H	66-68
1.099	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-CH <sub>3</sub>	H	H	resin
1.100	2-OCH <sub>3</sub> , 4-F	2-CH <sub>3</sub>	H	H	resin
1.101	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-CN	H	H	crystalline
1.102	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-OCH <sub>3</sub>	H	H	resin
1.103	2-OCH <sub>3</sub> , 4-F	3-OCH <sub>3</sub>	H	H	resin
1.104	2-OCH <sub>3</sub> , 4-F	2-CN	H	H	oil

Table 2: Compounds of formula I<sub>2</sub>:


Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
2.001	2-OCH <sub>3</sub> , 4-CN	2-F	H	H	132-134
2.002	2-F, 4-Cl	2-F	H	H	-
2.003	2-Cl, 4-Cl	2-F	H	H	-
2.004	2-OCH <sub>3</sub> , 4-F	2-F	H	H	resin
2.005	2-OCH <sub>3</sub> , 4-Cl	2-F	H	H	-
2.006	2-OCH <sub>3</sub> , 4-Br	2-F	H	H	-
2.007	2-CF <sub>3</sub> , 4-F	2-F	H	H	-
2.008	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-F	H	H	-
2.009	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-F	H	H	-
2.010	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-F	H	H	amorphous
2.011	2-OCH <sub>3</sub> , 4-F	H	H	H	crystalline
2.012	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	H	H	H	crystalline
2.013	2-OCH <sub>3</sub> , 4-CN	2-OCH <sub>3</sub>	H	H	-
2.014	2-F, 4-Cl	2-OCH <sub>3</sub>	H	H	-
2.015	2-Cl, 4-Cl	2-OCH <sub>3</sub>	H	H	-

- 36 -

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
2.016	2-OCH <sub>3</sub> , 4-F	2-OCH <sub>3</sub>	H	H	-
2.017	2-OCH <sub>3</sub> , 4-Cl	2-OCH <sub>3</sub>	H	H	-
2.018	2-OCH <sub>3</sub> , 4-Br	2-OCH <sub>3</sub>	H	H	-
2.019	2-CF <sub>3</sub> , 4-F	2-OCH <sub>3</sub>	H	H	-
2.020	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-OCH <sub>3</sub>	H	H	-
2.021	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-OCH <sub>3</sub>	H	H	-
2.022	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-OCH <sub>3</sub>	H	H	-
2.023	2-OCH <sub>3</sub> , 4-F	2-OCH <sub>3</sub> , 5-NH <sub>2</sub>	H	H	amorphous
2.024	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-OCH <sub>3</sub> , 5-NH <sub>2</sub>	H	H	amorphous
2.025	2-OCH <sub>3</sub> , 4-F	2-OCH <sub>3</sub> , 5-NHC(O)O-t-C <sub>4</sub> H <sub>9</sub>	H	H	oil
2.026	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-OCH <sub>3</sub> , 5-NHC(O)O-t-C <sub>4</sub> H <sub>9</sub>	H	H	crystalline
2.027	2-OCH <sub>3</sub> , 4-CN	2-Cl	H	H	-
2.028	2-F, 4-Cl	2-Cl	H	H	-
2.029	2-Cl, 4-Cl	2-Cl	H	H	-
2.030	2-OCH <sub>3</sub> , 4-F	2-Cl	H	H	-
2.031	2-OCH <sub>3</sub> , 4-Cl	2-Cl	H	H	-
2.032	2-OCH <sub>3</sub> , 4-Br	2-Cl	H	H	-
2.033	2-CF <sub>3</sub> , 4-F	2-Cl	H	H	-
2.034	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-Cl	H	H	-
2.035	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-Cl	H	H	-
2.036	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-Cl	H	H	-
2.037	2-OCH <sub>3</sub> , 4-CN	2-CH <sub>2</sub> CN	H	H	-
2.038	2-F, 4-Cl	2-CH <sub>2</sub> CN	H	H	-
2.039	2-Cl, 4-Cl	2-CH <sub>2</sub> CN	H	H	-
2.040	2-OCH <sub>3</sub> , 4-F	2-CH <sub>2</sub> CN	H	H	83-84
2.041	2-OCH <sub>3</sub> , 4-Cl	2-CH <sub>2</sub> CN	H	H	-
2.042	2-OCH <sub>3</sub> , 4-Br	2-CH <sub>2</sub> CN	H	H	-
2.043	2-CF <sub>3</sub> , 4-F	2-CH <sub>2</sub> CN	H	H	-
2.044	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-CH <sub>2</sub> CN	H	H	-
2.045	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-CH <sub>2</sub> CN	H	H	-
2.046	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-CH <sub>2</sub> CN	H	H	resin
2.047	2-OCH <sub>3</sub> , 4-CN	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	142-144

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
2.048	2-F, 4-Cl	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.049	2-Cl, 4-Cl	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.050	2-OCH <sub>3</sub> , 4-F	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.051	2-OCH <sub>3</sub> , 4-Cl	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.052	2-OCH <sub>3</sub> , 4-Br	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.053	2-CF <sub>3</sub> , 4-F	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.054	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.055	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.056	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-
2.057	2-OCH <sub>3</sub> , 4-CN	2-CH(CH <sub>3</sub> )CN	H	H	-
2.058	2-F, 4-Cl	2-CH(CH <sub>3</sub> )CN	H	H	-
2.059	2-Cl, 4-Cl	2-CH(CH <sub>3</sub> )CN	H	H	-
2.060	2-OCH <sub>3</sub> , 4-F	2-CH(CH <sub>3</sub> )CN	H	H	-
2.061	2-OCH <sub>3</sub> , 4-Cl	2-CH(CH <sub>3</sub> )CN	H	H	-
2.062	2-OCH <sub>3</sub> , 4-Br	2-CH(CH <sub>3</sub> )CN	H	H	-
2.063	2-CF <sub>3</sub> , 4-F	2-CH(CH <sub>3</sub> )CN	H	H	-
2.064	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-CH(CH <sub>3</sub> )CN	H	H	-
2.065	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	2-CH(CH <sub>3</sub> )CN	H	H	-
2.066	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-CH(CH <sub>3</sub> )CN	H	H	-
2.067	2-OCH <sub>3</sub> , 4-F	2-Cl	CH <sub>3</sub>	H	-
2.068	2-OCH <sub>3</sub> , 4-Cl	2-Cl	CH <sub>3</sub>	H	-
2.069	2-OCH <sub>3</sub> , 4-Br	2-Cl	CH <sub>3</sub>	H	-
2.070	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-Cl	CH <sub>3</sub>	H	-
2.071	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-Cl	CH <sub>3</sub>	H	-
2.072	2-OCH <sub>3</sub> , 4-F	2-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
2.073	2-OCH <sub>3</sub> , 4-Cl	2-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
2.074	2-OCH <sub>3</sub> , 4-Br	2-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
2.075	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
2.076	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-CH <sub>2</sub> CN	CH <sub>3</sub>	CH <sub>3</sub>	-
2.077	2-OCH <sub>3</sub> , 4-F	2-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.078	2-OCH <sub>3</sub> , 4-Cl	2-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.079	2-OCH <sub>3</sub> , 4-Br	2-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.080	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	2-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.081	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	2-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
2.082	2-OCH <sub>3</sub> , 4-F	3-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.083	2-OCH <sub>3</sub> , 4-Cl	3-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.084	2-OCH <sub>3</sub> , 4-Br	3-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.085	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	3-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.086	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-CH <sub>2</sub> CN	CH <sub>3</sub>	H	-
2.087	2-OCH <sub>3</sub>	2-CH <sub>2</sub> CN	H	H	-
2.088	2-OCH <sub>3</sub>	3-CH <sub>2</sub> CN	H	H	-
2.089	2-OCH <sub>3</sub>	2-F	H	H	-
2.090	2-OCH <sub>3</sub>	2-Cl	H	H	-
2.091	2-OCH <sub>3</sub>	2-Br	H	H	-

Table 3: Compounds of formula I<sub>3</sub>:

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
3.001	2-OCH <sub>3</sub> , 4-CN	4-CH <sub>3</sub>	H	H	-
3.002	2-F, 4-Cl	4-CH <sub>3</sub>	H	H	-
3.003	2-Cl, 4-Cl	4-CH <sub>3</sub>	H	H	-
3.004	2-OCH <sub>3</sub> , 4-F	4-CH <sub>3</sub>	H	H	crystalline
3.005	2-OCH <sub>3</sub> , 4-Cl	4-CH <sub>3</sub>	H	H	-
3.006	2-OCH <sub>3</sub> , 4-Br	4-CH <sub>3</sub>	H	H	-
3.007	2-CF <sub>3</sub> , 4-F	4-CH <sub>3</sub>	H	H	-
3.008	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	4-CH <sub>3</sub>	H	H	-
3.009	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	4-CH <sub>3</sub>	H	H	-
3.010	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	4-CH <sub>3</sub>	H	H	
3.011	H	4-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.012	2-OCH <sub>3</sub> , 4-CH <sub>2</sub> CN	4-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.013	4-NO <sub>2</sub>	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.014	2-OCH <sub>3</sub>	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
3.015	4-CH <sub>2</sub> CN	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.016	2-OCH <sub>3</sub> , 4-CH <sub>2</sub> CN	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.017	4-CN	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.018	4-CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.019	2-Cl, 6-Cl	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.020	H	3-OH, 6-CH <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.021	2-OCH <sub>3</sub> , 4-F	6-CH <sub>3</sub>	H	H	oil
3.022	2-OCH <sub>3</sub> , 4-F	5-CH <sub>3</sub>	H	H	oil
3.023	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	5-CH <sub>3</sub>	H	H	crystalline
3.024	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	6-CH <sub>3</sub>	H	H	crystalline
3.025	4-OC <sub>6</sub> H <sub>5</sub>	H	H	H	-
3.026	2-OCH <sub>3</sub> , 4-CH <sub>2</sub> CN	H	H	H	MS: [M+H] <sup>+</sup>
3.027	4-CH <sub>2</sub> CN	H	H	H	MS: [M+H] <sup>+</sup>
3.028	H	H	H	H	MS: [M+H] <sup>+</sup>
3.029	2-OCH <sub>3</sub> , 4-CN	5-CF <sub>3</sub>	H	H	94-95
3.030	2-F, 4-Cl	5-CF <sub>3</sub>	H	H	-
3.031	2-OCH <sub>3</sub> , 4-F	5-CF <sub>3</sub>	H	H	crystalline
3.032	2-OCH <sub>3</sub> , 4-Cl	5-CF <sub>3</sub>	H	H	-
3.033	2-OCH <sub>3</sub> , 4-Br	5-CF <sub>3</sub>	H	H	-
3.034	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	5-CF <sub>3</sub>	H	H	-
3.035	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	5-CF <sub>3</sub>	H	H	-
3.036	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	5-CF <sub>3</sub>	H	H	crystalline
3.037	4-CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	5-CF <sub>3</sub>	H	H	MS: [M+H] <sup>+</sup>
3.038	2-OCH <sub>3</sub> , 4-CN	4-CH <sub>2</sub> CN	H	H	-
3.039	2-F, 4-Cl	4-CH <sub>2</sub> CN	H	H	-
3.040	2-Cl, 4-Cl	4-CH <sub>2</sub> CN	H	H	-
3.041	2-OCH <sub>3</sub> , 4-F	4-CH <sub>2</sub> CN	H	H	-
3.042	2-OCH <sub>3</sub> , 4-Cl	4-CH <sub>2</sub> CN	H	H	-
3.043	2-OCH <sub>3</sub> , 4-Br	4-CH <sub>2</sub> CN	H	H	-
3.044	2-CF <sub>3</sub> , 4-F	4-CH <sub>2</sub> CN	H	H	-
3.045	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	4-CH <sub>2</sub> CN	H	H	-
3.046	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	4-CH <sub>2</sub> CN	H	H	-
3.047	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	4-CH <sub>2</sub> CN	H	H	-
3.048	2-OCH <sub>3</sub>	4-CH <sub>2</sub> CN	H	H	-



Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
3.049	2-OCH <sub>3</sub>	4-Cl	H	H	-
3.050	2-OCH <sub>3</sub>	4-Br	H	H	-
3.051	2-OCH <sub>3</sub>	6-CH <sub>2</sub> CN	H	H	106
3.052	2-OCH <sub>3</sub>	6-Cl	H	H	-
3.053	2-OCH <sub>3</sub>	6-Br	H	H	-
3.054	2-OCH <sub>3</sub> , 4-CN	5-Cl	H	H	-
3.055	2-F, 4-Cl	5-Cl	H	H	-
3.056	2-OCH <sub>3</sub> , 4-F	5-Cl	H	H	-
3.057	2-OCH <sub>3</sub> , 4-Cl	5-Cl	H	H	-
3.058	2-OCH <sub>3</sub> , 4-Br	5-Cl	H	H	-
3.059	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	5-Cl	H	H	-
3.060	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	5-Cl	H	H	-
3.061	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	5-Cl	H	H	-
3.062	4-OCH <sub>2</sub> CH <sub>2</sub> N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	5-Cl	H	H	58-60
3.063	2-OCH <sub>3</sub> , 4-CN	6-Br	H	H	84-85
3.064	2-F, 4-Cl	6-Br	H	H	-
3.065	2-Cl, 4-Cl	6-Br	H	H	-
3.066	2-OCH <sub>3</sub> , 4-F	6-Br	H	H	crystalline
3.067	2-OCH <sub>3</sub> , 4-Cl	6-Br	H	H	-
3.068	2-OCH <sub>3</sub> , 4-Br	6-Br	H	H	-
3.069	2-CF <sub>3</sub> , 4-F	6-Br	H	H	-
3.070	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	6-Br	H	H	-
3.071	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	6-Br	H	H	-
3.072	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	6-Br	H	H	crystalline
3.073	2-OCH <sub>3</sub> , 4-F	4-CH <sub>3</sub>	CH <sub>3</sub>	H	-
3.074	2-OCH <sub>3</sub> , 4-Cl	4-CH <sub>3</sub>	CH <sub>3</sub>	H	-
3.075	2-OCH <sub>3</sub> , 4-Br	4-CH <sub>3</sub>	CH <sub>3</sub>	H	-
3.076	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	4-CH <sub>3</sub>	CH <sub>3</sub>	H	-
3.077	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	4-CH <sub>3</sub>	CH <sub>3</sub>	H	-
3.078	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	4-CH <sub>3</sub>	CH <sub>3</sub>	H	-
3.079	2-OCH <sub>3</sub> , 4-F	4-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	-
3.080	2-OCH <sub>3</sub> , 4-Cl	4-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	-
3.081	2-OCH <sub>3</sub> , 4-Br	4-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	-
3.082	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	4-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	-

Comp. No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	Phys. data m.p. (°C)
3.083	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	4-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	-
3.084	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	4-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	-
3.085	2-OCH <sub>3</sub> , 4-F	3-OH	H	H	crystalline
3.086	2-OCH <sub>3</sub> , 4-Cl	3-OH	H	H	-
3.087	2-OCH <sub>3</sub> , 4-Br	3-OH	H	H	-
3.088	2-OCH <sub>3</sub> , 4-CF <sub>3</sub>	3-OH	H	H	-
3.089	2-OCH <sub>3</sub> , 4-CH <sub>3</sub>	3-OH	H	H	-
3.090	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	3-OH	H	H	crystalline
3.091	4-CH <sub>2</sub> CN	3-OC <sub>2</sub> H <sub>5</sub>	H	H	MS: [M+H] <sup>+</sup>
3.092	2-OCH <sub>3</sub>	3-OC <sub>2</sub> H <sub>5</sub>	H	H	MS: [M+H] <sup>+</sup>
3.093	2-OCH <sub>3</sub> , 4-CH <sub>2</sub> CN	3-OC <sub>2</sub> H <sub>5</sub>	H	H	MS: [M+H] <sup>+</sup>
3.094	2-OCH <sub>3</sub> , 4-CN	3-OC <sub>2</sub> H <sub>5</sub>	H	H	MS: [M+H] <sup>+</sup>
3.095	2-OCH <sub>3</sub> , 4-F	6-CH <sub>2</sub> CN	H	H	resin
3.096	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	6-CH <sub>2</sub> CN	H	H	solid
3.097	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	5-CH <sub>2</sub> CN	H	H	crystalline
3.098	2-OCH <sub>3</sub> , 4-F	5-CH <sub>2</sub> CN	H	H	resin
3.099	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	6-OCH <sub>3</sub>	H	H	resin
3.100	2-OCH <sub>3</sub> , 4-F	6-OCH <sub>3</sub>	H	H	resin
3.101	2-OCH <sub>3</sub> , 4-CH=NOCH <sub>3</sub>	H	H	H	resin
3.102	2-OCH <sub>3</sub> , 4-F	H	H	H	oil

### Biological Examples

#### Example B1: Herbicidal action prior to emergence of the plants (pre-emergence action)

Monocotyledonous and dicotyledonous test plants are sown in standard soil in pots. Immediately after sowing, the test compounds, in the form of an aqueous suspension (prepared from a wettable powder (Example F3, b) according to WO 97/34485) or in the form of an emulsion (prepared from an emulsifiable concentrate (Example F1, c) according to WO 97/34485), are applied by spraying in an optimum concentration (500 litres of water/ha). The test plants are then grown in a greenhouse under optimum conditions. After a test duration of 4 weeks, the test is evaluated in accordance with a scale of nine ratings (1 = total damage, 9 = no action). Ratings of from 1 to 4 (especially from 1 to 3) indicate good to very good herbicidal action.

Test plants: Panicum, Echinochloa (Ds), Amaranthus, Chenopodium, Stellaria, Veronica.

Table B1:

Concentration 1000 g of active ingredient/ha

Comp. No.	Panicum	Echinochloa (Ds)	Amaranthus	Chenopodium	Stellaria	Veronica
1.010	3	-	1	1	1	1
1.004	2	2	1	1	1	1
3.004	2	2	1	1	1	1

The same results are obtained when the compounds of formula I are formulated in accordance with the other Examples analogously to WO 97/34485.

Example B2: Post-emergence herbicidal action

Monocotyledonous and dicotyledonous test plants are sown in standard soil in pots. When the test plants are at the 2- to 3-leaf stage, the test compounds, in the form of an aqueous suspension (prepared from a wettable powder (Example F3, b) according to WO 97/34485) or in the form of an emulsion (prepared from an emulsifiable concentrate (Example F1, c) according to WO 97/34485), are applied by spraying in an optimum concentration (500 litres of water/ha). The test plants are then grown on in a greenhouse under optimum conditions. After a test duration of 2 to 3 weeks, the test is evaluated in accordance with a scale of nine ratings (1 = total damage, 9 = no action). Ratings of from 1 to 4 (especially from 1 to 3) indicate good to very good herbicidal action.

Test plants: Panicum, Euphorbia, Amaranthus, Chenopodium, Stellaria, Veronica.

Table B2:

Concentration 1000 g of active ingredient/ha


Comp. No.	Panicum	Euphorbia	Amaranthus	Chenopodium	Stellaria	Veronica
1.010	4	1	1	1	2	3
1.004	-	2	1	1	2	2
3.004	5	3	1	1	2	3

In the above Tables B1 and B2 " - " means that no data are available for that indication.

The same results are obtained when the compounds of formula I are formulated in accordance with the other Examples analogously to WO 97/34485.

What is claimed is:

## 1. A compound of formula I



wherein

Z is =N- or  $\text{—}\overset{\text{||}}{\text{N}}\text{—}\overset{+}{\text{O}}\text{—}$  ;

n is 0, 1, 2, 3, 4 or 5;

each  $R_1$  independently of any others is halogen, -CN, -SCN, -SF<sub>5</sub>, -NO<sub>2</sub>, -NR<sub>5</sub>R<sub>6</sub>, -CO<sub>2</sub>R<sub>7</sub>, -CONR<sub>8</sub>R<sub>9</sub>, -C(R<sub>10</sub>)=NOR<sub>11</sub>, -COR<sub>12</sub>, -OR<sub>13</sub>, -SR<sub>14</sub>, -SOR<sub>15</sub>, -SO<sub>2</sub>R<sub>16</sub>, -OSO<sub>2</sub>R<sub>17</sub>, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl; or is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl or C<sub>2</sub>-C<sub>8</sub>alkynyl substituted by one or more halogen, -CN, -NO<sub>2</sub>, -NR<sub>18</sub>R<sub>19</sub>, -CO<sub>2</sub>R<sub>20</sub>, -CONR<sub>21</sub>R<sub>22</sub>, -COR<sub>23</sub>, -C(R<sub>24</sub>)=NOR<sub>25</sub>, -C(S)NR<sub>26</sub>R<sub>27</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>28</sub>, -OR<sub>29</sub>, -SR<sub>30</sub>, -SOR<sub>31</sub>, -SO<sub>2</sub>R<sub>32</sub> or C<sub>3</sub>-C<sub>6</sub>cycloalkyl substituents; or

each  $R_1$  independently of any others is C<sub>3</sub>-C<sub>6</sub>cycloalkyl substituted by one or more halogen, -CN, -NO<sub>2</sub>, -NR<sub>18</sub>R<sub>19</sub>, -CO<sub>2</sub>R<sub>20</sub>, -CONR<sub>21</sub>R<sub>22</sub>, -COR<sub>23</sub>, -C(R<sub>24</sub>)=NOR<sub>25</sub>, -C(S)NR<sub>26</sub>R<sub>27</sub>,

-C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>28</sub>, -SR<sub>30</sub>, -SOR<sub>31</sub>, -SO<sub>2</sub>R<sub>32</sub> or C<sub>3</sub>-C<sub>6</sub>cycloalkyl substituents; or

each  $R_1$  independently of any others is phenyl, which may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

two adjacent  $R_1$  together form a C<sub>1</sub>-C<sub>7</sub>alkylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by C<sub>1</sub>-C<sub>6</sub>alkyl or C<sub>1</sub>-C<sub>6</sub>alkoxy, the total number of ring atoms being at least 5 and at most 9; or

two adjacent  $R_1$  together form a C<sub>2</sub>-C<sub>7</sub>alkenylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by C<sub>1</sub>-C<sub>6</sub>alkyl or C<sub>1</sub>-C<sub>6</sub>alkoxy, the total number of ring atoms being at least 5 and at most 9;

$R_3$  and  $R_4$  are each independently of the other hydrogen, halogen, -CN, C<sub>1</sub>-C<sub>4</sub>alkyl or C<sub>1</sub>-C<sub>4</sub>alkoxy; or

$R_3$  and  $R_4$  together are C<sub>2</sub>-C<sub>5</sub>alkylene;

$R_5$  is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>6</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl; wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>5</sub> and R<sub>6</sub> together are a C<sub>2</sub>-C<sub>5</sub>alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

R<sub>7</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>8</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>9</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents, or

R<sub>9</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>8</sub> and R<sub>9</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>10</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>11</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>12</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>13</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl; or

R<sub>13</sub> is phenyl or phenyl-C<sub>1</sub>-C<sub>6</sub>alkyl, wherein both phenyl rings may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>8</sub>alkylthio, C<sub>1</sub>-C<sub>8</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>8</sub>alkylsulfonyl substituents, or

R<sub>13</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)-amino or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>14</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>15</sub>, R<sub>16</sub> and R<sub>17</sub> are each independently of the others C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>18</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>19</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>18</sub> and R<sub>19</sub> together are a C<sub>2</sub>-C<sub>5</sub>alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

R<sub>20</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>21</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>22</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents, or

R<sub>22</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>21</sub> and R<sub>22</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>23</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>24</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>25</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>26</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>27</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents, or

R<sub>27</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>26</sub> and R<sub>27</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>28</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>29</sub> and R<sub>30</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>31</sub> and R<sub>32</sub> are each independently of the other C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

m is 0, 1, 2, 3 or 4;

each R<sub>2</sub> independently of any others is halogen, -CN, -SCN, -OCN, -N<sub>3</sub>, -SF<sub>5</sub>, -NO<sub>2</sub>, -NR<sub>33</sub>R<sub>34</sub>, -CO<sub>2</sub>R<sub>35</sub>, -CONR<sub>36</sub>R<sub>37</sub>, -C(R<sub>38</sub>)=NOR<sub>39</sub>, -COR<sub>40</sub>, -OR<sub>41</sub>, -SR<sub>42</sub>, -SOR<sub>43</sub>, -SO<sub>2</sub>R<sub>44</sub>, -OSO<sub>2</sub>R<sub>45</sub>, -N([CO]<sub>p</sub>R<sub>46</sub>)COR<sub>47</sub>, -N(OR<sub>54</sub>)COR<sub>55</sub>, -N(R<sub>56</sub>)SO<sub>2</sub>R<sub>57</sub>, -N(SO<sub>2</sub>R<sub>58</sub>)SO<sub>2</sub>R<sub>59</sub>, -N=C(OR<sub>60</sub>)R<sub>61</sub>, -CR<sub>62</sub>(OR<sub>63</sub>)OR<sub>64</sub>, -OC(O)NR<sub>65</sub>R<sub>66</sub>, -SC(O)NR<sub>67</sub>R<sub>68</sub>, -OC(S)NR<sub>69</sub>R<sub>70</sub> or -N-phthalimide; or

R<sub>2</sub> is a 5- to 7-membered heterocyclic ring system which may be aromatic or partially or fully saturated and may contain from 1 to 4 hetero atoms selected from nitrogen, oxygen and sulfur, it being possible for that heterocyclic ring system in turn to be substituted by one or

more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, hydroxy-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>4</sub>alkyl, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl substituents; R<sub>33</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl; and

R<sub>34</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or R<sub>33</sub> and R<sub>34</sub> together are a C<sub>2</sub>-C<sub>5</sub>alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

R<sub>35</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>36</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>37</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents, or

R<sub>37</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>36</sub> and R<sub>37</sub> together are C<sub>3</sub>-C<sub>5</sub>alkylene;

R<sub>38</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>39</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>40</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>8</sub>alkylthio, -C(O)-C(O)OC<sub>1</sub>-C<sub>4</sub>alkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>41</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>8</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl, C<sub>3</sub>-C<sub>8</sub>alkenyloxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylthio-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl-C<sub>1</sub>-C<sub>6</sub>alkyl or C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl-C<sub>1</sub>-C<sub>6</sub>alkyl; or R<sub>41</sub> is phenyl or phenyl-C<sub>1</sub>-C<sub>6</sub>alkyl, wherein both phenyl rings may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, or -S(O)<sub>2</sub>C<sub>1</sub>-C<sub>8</sub>alkyl substituents, or

R<sub>41</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)amino or -CN substituents;

R<sub>42</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>43</sub> and R<sub>44</sub> are each independently of the other C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>45</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or

R<sub>45</sub> is phenyl, it being possible for the phenyl ring to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>8</sub>alkylthio, C<sub>1</sub>-C<sub>8</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>8</sub>alkylsulfonyl substituents;

R<sub>46</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or C<sub>1</sub>-C<sub>4</sub>haloalkyl;

R<sub>47</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more halogen, -CN, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, -NR<sub>48</sub>COR<sub>49</sub>, -NR<sub>50</sub>SO<sub>2</sub>R<sub>51</sub> or -NR<sub>52</sub>CO<sub>2</sub>R<sub>53</sub> substituents, or R<sub>47</sub> is phenyl or benzyl, each of which may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

p is 0 or 1;

R<sub>48</sub>, R<sub>49</sub>, R<sub>50</sub>, R<sub>51</sub>, R<sub>52</sub> and R<sub>53</sub> are each independently of the others hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, phenyl, benzyl or naphthyl, it being possible for the three last-mentioned aromatic radicals in turn to be substituted by one or more halogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino, -NH<sub>2</sub>, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>54</sub> and R<sub>55</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl or phenyl, whereby the phenyl ring may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>8</sub>alkylthio, C<sub>1</sub>-C<sub>8</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>8</sub>alkylsulfonyl substituents;

R<sub>56</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or benzyl, it being possible for benzyl in turn to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>8</sub>alkylthio, C<sub>1</sub>-C<sub>8</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>8</sub>alkylsulfonyl substituents;

R<sub>57</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, phenyl, benzyl or naphthyl, it being possible for the three last-mentioned aromatic rings to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino, -NH<sub>2</sub>, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>58</sub> and R<sub>59</sub> are each independently of the other C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl, benzyl or naphthyl, it being possible for the three last-mentioned aromatic rings to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino, -NH<sub>2</sub>, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>60</sub> and R<sub>61</sub> are each independently of the other hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>62</sub>, R<sub>63</sub> and R<sub>64</sub> are each independently of the others hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or



$R_{63}$  and  $R_{64}$  together form a  $C_2$ - $C_5$ alkylene bridge;

$R_{65}$ ,  $R_{66}$ ,  $R_{67}$ ,  $R_{68}$ ,  $R_{69}$  and  $R_{70}$  are each independently of the others hydrogen or  $C_1$ - $C_8$ alkyl, or

$R_{65}$  and  $R_{66}$  together or  $R_{67}$  and  $R_{68}$  together or  $R_{69}$  and  $R_{70}$  together form a  $C_2$ - $C_5$ alkylene bridge; or

each  $R_2$  independently of any others is  $C_1$ - $C_8$ alkyl, or is  $C_1$ - $C_8$ alkyl mono- or poly-substituted by halogen, -CN, -N<sub>3</sub>, -SCN, -NO<sub>2</sub>, -NR<sub>71</sub>R<sub>72</sub>, -CO<sub>2</sub>R<sub>73</sub>, -CONR<sub>74</sub>R<sub>75</sub>, -COR<sub>76</sub>, -C(R<sub>77</sub>)=NOR<sub>78</sub>, -C(S)NR<sub>79</sub>R<sub>80</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>81</sub>, -OR<sub>82</sub>, -SR<sub>83</sub>, -SOR<sub>84</sub>, -SO<sub>2</sub>R<sub>85</sub>, -O(SO<sub>2</sub>)R<sub>86</sub>,

-N(R<sub>87</sub>)CO<sub>2</sub>R<sub>88</sub>, -N(R<sub>89</sub>)COR<sub>90</sub>, -S<sup>+</sup>(R<sub>91</sub>)<sub>2</sub>, -N<sup>+</sup>(R<sub>92</sub>)<sub>3</sub>, -Si(R<sub>93</sub>)<sub>3</sub> or  $C_3$ - $C_6$ cycloalkyl; or

each  $R_2$  independently of any others is  $C_1$ - $C_8$ alkyl substituted by a 5- to 7-membered heterocyclic ring system, which may be aromatic or partially or fully saturated and may contain from 1 to 4 hetero atoms selected from nitrogen, oxygen and sulfur, it being possible for that heterocyclic ring system in turn to be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl, hydroxy- $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_4$ alkoxy- $C_1$ - $C_4$ alkyl, -CN, -NO<sub>2</sub>,  $C_1$ - $C_6$ alkylthio,  $C_1$ - $C_6$ alkylsulfinyl or  $C_1$ - $C_6$ alkylsulfonyl substituents; or

each  $R_2$  independently of any others is  $C_2$ - $C_8$ alkenyl, or is  $C_2$ - $C_8$ alkenyl mono- or poly-substituted by halogen, -CN, -NO<sub>2</sub>, -CO<sub>2</sub>R<sub>94</sub>, -CONR<sub>95</sub>R<sub>96</sub>, -COR<sub>97</sub>, -C(R<sub>98</sub>)=NOR<sub>99</sub>, -C(S)NR<sub>100</sub>R<sub>101</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>102</sub>, -OR<sub>103</sub>, -Si(R<sub>104</sub>)<sub>3</sub> or  $C_3$ - $C_6$ cycloalkyl; or

each  $R_2$  independently of any others is  $C_2$ - $C_8$ alkynyl, or is  $C_2$ - $C_8$ alkynyl mono- or poly-substituted by halogen, -CN, -CO<sub>2</sub>R<sub>105</sub>, -CONR<sub>106</sub>R<sub>107</sub>, -COR<sub>108</sub>, -C(R<sub>109</sub>)=NOR<sub>110</sub>, -C(S)NR<sub>111</sub>R<sub>112</sub>, -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>113</sub>, -OR<sub>114</sub>, -Si(R<sub>115</sub>)<sub>3</sub> or  $C_3$ - $C_6$ cycloalkyl; or

each  $R_2$  independently of any others is  $C_3$ - $C_6$ cycloalkyl, or is  $C_3$ - $C_6$ cycloalkyl mono- or poly-substituted by halogen, -CN, -CO<sub>2</sub>R<sub>116</sub>, -CONR<sub>117</sub>R<sub>118</sub>, -COR<sub>119</sub>, -C(R<sub>120</sub>)=NOR<sub>121</sub>, -C(S)NR<sub>122</sub>R<sub>123</sub> or -C(C<sub>1</sub>-C<sub>4</sub>alkylthio)=NR<sub>124</sub>; or

two adjacent  $R_2$  together form a  $C_1$ - $C_7$ alkylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9; or

two adjacent  $R_2$  together form a  $C_2$ - $C_7$ alkenylene bridge, which may be interrupted by 1 or 2 non-adjacent oxygen atoms and may be substituted by  $C_1$ - $C_6$ alkyl or  $C_1$ - $C_6$ alkoxy, the total number of ring atoms being at least 5 and at most 9;

$R_{71}$  is hydrogen or  $C_1$ - $C_8$ alkyl;

$R_{72}$  is hydrogen,  $C_1$ - $C_8$ alkyl,  $C_3$ - $C_6$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or

R<sub>71</sub> and R<sub>72</sub> together are a C<sub>2</sub>-C<sub>5</sub>alkylene chain, which may be interrupted by an oxygen or a sulfur atom;

R<sub>73</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, it being possible for phenyl in turn to be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>74</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>75</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl or C<sub>3</sub>-C<sub>7</sub>cycloalkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxy or -CN substituents; or

R<sub>75</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>74</sub> and R<sub>75</sub> together are a C<sub>2</sub>-C<sub>5</sub>alkylene chain, which may be interrupted by an oxygen or sulfur atom;

R<sub>76</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>77</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>78</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl; and

R<sub>79</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>80</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents; or

R<sub>80</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>79</sub> and R<sub>80</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>81</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>82</sub> is -Si(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>3</sub>, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or C<sub>1</sub>-C<sub>8</sub>alkyl, whereby C<sub>1</sub>-C<sub>8</sub>alkyl is mono- or poly-substituted by halogen, -CN, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)amino or C<sub>1</sub>-C<sub>4</sub>alkoxy;

R<sub>83</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl or C<sub>1</sub>-C<sub>8</sub>alkyl, whereby C<sub>1</sub>-C<sub>8</sub>alkyl is mono- or poly-substituted by halogen, -CN, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylamino, di(C<sub>1</sub>-C<sub>6</sub>alkyl)amino or C<sub>1</sub>-C<sub>4</sub>alkoxy;

R<sub>84</sub>, R<sub>85</sub> and R<sub>86</sub> are each independently of the others C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, or C<sub>1</sub>-C<sub>8</sub>alkyl which is substituted by one or more halogen, -CN or C<sub>1</sub>-C<sub>4</sub>alkoxy substituents;

R<sub>87</sub> and R<sub>89</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl or C<sub>1</sub>-C<sub>8</sub>alkoxy;  
R<sub>88</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl;  
R<sub>90</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;  
R<sub>91</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl;  
R<sub>92</sub> and R<sub>93</sub> are each independently of the other C<sub>1</sub>-C<sub>6</sub>alkyl;  
R<sub>94</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, each of which may be mono- or poly-substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;  
R<sub>95</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;  
R<sub>96</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl or -CN substituents; or  
R<sub>96</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or  
R<sub>95</sub> and R<sub>96</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;  
R<sub>97</sub> and R<sub>98</sub> are each independently of the other hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;  
R<sub>99</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;  
R<sub>100</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;  
R<sub>101</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl or -CN substituents; or  
R<sub>101</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or  
R<sub>100</sub> and R<sub>101</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;  
R<sub>102</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;  
R<sub>103</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, -Si(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>3</sub>, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl;  
R<sub>104</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl;  
R<sub>105</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, each of which may be mono- or poly-substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;  
R<sub>106</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;  
R<sub>107</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents; or

R<sub>107</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>106</sub> and R<sub>107</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>108</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>109</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>110</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>111</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>112</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl or -CN substituents; or

R<sub>112</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>111</sub> and R<sub>112</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>113</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>114</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, -Si(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>3</sub>, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl;

R<sub>115</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl;

R<sub>116</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl or C<sub>3</sub>-C<sub>8</sub>alkynyl, each of which may be mono- or poly-substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkoxy or phenyl substituents, wherein phenyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents;

R<sub>117</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>118</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl or -CN substituents; or

R<sub>118</sub> is C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, -CN, -NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylsulfinyl or C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl substituents; or

R<sub>117</sub> and R<sub>118</sub> together are C<sub>2</sub>-C<sub>5</sub>alkylene;

R<sub>119</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>120</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkyl;

R<sub>121</sub> is hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>alkenyl, C<sub>3</sub>-C<sub>8</sub>alkynyl, C<sub>1</sub>-C<sub>4</sub>haloalkyl or C<sub>3</sub>-C<sub>6</sub>haloalkenyl;

R<sub>122</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

R<sub>123</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl, or is C<sub>1</sub>-C<sub>8</sub>alkyl substituted by one or more -COOH, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl or -CN substituents; or


$R_{123}$  is  $C_3$ - $C_8$ alkenyl,  $C_3$ - $C_8$ alkynyl, phenyl or benzyl, wherein phenyl and benzyl may in turn be substituted by one or more halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ haloalkyl,  $C_1$ - $C_4$ alkoxy, -CN, -NO<sub>2</sub>,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylsulfinyl or  $C_1$ - $C_4$ alkylsulfonyl substituents; or

$R_{122}$  and  $R_{123}$  together are  $C_2$ - $C_5$ alkylene; and


$R_{124}$  is hydrogen or  $C_1$ - $C_8$ alkyl,

or an agrochemically acceptable salt or any stereoisomer or tautomer of a compound of formula I.


2. A process for the preparation of a compound of formula I according to claim 1, which process comprises reacting a compound of formula II




wherein  $R_1$  and  $n$  are as defined in claim 1, in the presence of a base, with a compound of formula III




wherein  $R_3$  and  $R_4$  are as defined in claim 1 and  $X_1$  is O-tosyl, O-mesyl, chlorine, bromine or iodine, to form a compound of formula IV



wherein  $R_1$ ,  $R_3$ ,  $R_4$  and  $n$  are as defined, and then coupling that compound with a compound of formula V or Va



wherein  $R_2$  and  $m$  are as defined in claim 1 and  $A$  is a leaving group, in the presence of a palladium catalyst, and, if desired, oxidising the resulting pyridine derivative of formula I wherein  $Z$  is  $=N-$  to form the corresponding pyridine N-oxide of formula I wherein  $Z$  is



3. A herbicidal and plant-growth-inhibiting composition, comprising a herbicidally effective amount of a compound of formula I on an inert carrier.
4. A method of controlling undesired plant growth, which method comprises applying a compound of formula I, or a composition comprising such a compound, in a herbicidally effective amount to plants or to the locus thereof.
5. A method of inhibiting plant growth, which method comprises applying a compound of formula I, or a composition comprising such a compound, in a herbicidally effective amount to plants or to the locus thereof.
6. A compound according to claim 1, wherein  $Z$  is  $=N-$ ; and each  $R_2$  independently of any others is  $C_2$ - $C_8$ alkenyl, or is  $C_2$ - $C_8$ alkenyl mono- or poly-substituted by  $-\text{CN}$ ,  $-\text{NO}_2$ ,  $-\text{CO}_2\text{R}_{94}$ ,  $-\text{CONR}_{95}\text{R}_{96}$ ,  $-\text{COR}_{97}$ ,  $-\text{C}(\text{R}_{98})=\text{NOR}_{99}$ ,  $-\text{C}(\text{S})\text{NR}_{100}\text{R}_{101}$ ,  $-\text{C}(\text{C}_1\text{-C}_4\text{alkylthio})=\text{NR}_{102}$ ,  $-\text{OR}_{103}$ ,  $-\text{Si}(\text{R}_{104})_3$  or  $\text{C}_3$ - $\text{C}_6$ cycloalkyl.
7. A compound according to claim 1, wherein each  $R_2$  independently of any others is halogen,  $-\text{CN}$ ,  $-\text{SCN}$ ,  $-\text{OCN}$ ,  $-\text{N}_3$ ,  $-\text{CONR}_{36}\text{R}_{37}$ ,  $-\text{C}(\text{R}_{38})=\text{NOR}_{39}$ ,  $-\text{COR}_{40}$ ,  $-\text{OR}_{41}$ ,  $-\text{SO}_2\text{R}_{45}$ ,  $-\text{N}([\text{CO}]_p\text{R}_{46})\text{COR}_{47}$ ,  $-\text{N}(\text{R}_{56})\text{SO}_2\text{R}_{57}$ ,  $-\text{N}(\text{SO}_2\text{R}_{58})\text{SO}_2\text{R}_{59}$ ,  $-\text{N}=\text{C}(\text{OR}_{60})\text{R}_{61}$  or  $\text{C}_1$ - $\text{C}_8$ alkyl, or is  $\text{C}_1$ - $\text{C}_8$ alkyl mono- or poly-substituted by halogen,  $-\text{CN}$ ,  $-\text{N}_3$ ,  $-\text{SCN}$ ,  $-\text{CONR}_{74}\text{R}_{75}$ ,  $-\text{COR}_{76}$ ,  $-\text{C}(\text{R}_{77})=\text{NOR}_{78}$ ,  $-\text{C}(\text{S})\text{NR}_{79}\text{R}_{80}$ ,  $-\text{OR}_{82}$ ,  $-\text{SOR}_{84}$ ,  $-\text{SO}_2\text{R}_{85}$  or  $-\text{N}(\text{R}_{89})\text{COR}_{90}$ .
8. A compound according to claim 1, wherein each  $R_1$  independently of any others is halogen,  $-\text{CN}$ ,  $\text{C}_1$ - $\text{C}_3$ alkyl,  $\text{C}_1$ - $\text{C}_3$ haloalkyl,  $\text{C}_1$ - $\text{C}_3$ cyanoalkyl,  $-\text{OR}_{13}$  or  $-\text{C}(\text{R}_{24})=\text{NOR}_{25}$ ;  $\text{R}_{13}$  is  $\text{C}_1$ - $\text{C}_3$ alkyl or  $\text{di}(\text{C}_1\text{-C}_4\text{-alkyl})\text{amino-C}_1\text{-C}_4\text{alkyl}$ ;  $\text{R}_{24}$  is hydrogen or methyl; and  $\text{R}_{25}$  is hydrogen or  $\text{C}_1$ - $\text{C}_3$ alkyl.
9. A compound according to claim 1, wherein  $\text{R}_3$  and  $\text{R}_4$  are each independently of the other hydrogen or methyl.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08878

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N43/40 C07D213/61 C07D213/64 C07D213/16 C07D213/73  
 C07D213/57 C07D213/75 C07D213/65 C07D213/74

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	WO 01 94339 A (EDMUNDS ANDREW ;LUETHY CHRISTOPH (CH); MESMAEKER ALAIN DE (CH); SY) 13 December 2001 (2001-12-13) claim 1	1-9
P,Y	WO 02 28182 A (HALL ROGER GRAHAM ;SCHAETZER JUERGEN (CH); EBERLE MARTIN (CH); REN) 11 April 2002 (2002-04-11) cited in the application claim 1	1-9
Y	WO 01 55066 A (EBERLE MARTIN ;ZELLER MARTIN (CH); EHRLER JUERG (CH); CRAIG GERALD) 2 August 2001 (2001-08-02) cited in the application claim 1	1-9
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the International filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the International filing date but later than the priority date claimed

- \*T\* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

31 October 2002

Date of mailing of the international search report

08/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Johnson, C

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08878

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PATENT ABSTRACTS OF JAPAN vol. 1999, no. 11, 30 September 1999 (1999-09-30) -& JP 11 147866 A (SANKYO CO LTD), 2 June 1999 (1999-06-02) cited in the application abstract; examples 1.1, 5.70, 5.71, 5.72 ----	1-9
X	DE 41 15 465 A (BEIERSDORF AG) 12 November 1992 (1992-11-12) formula (II), p. 7; examples 6'-12', 14', 15' ----	1,2,6-9
X	ATTWOOD M R ET AL: "TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 32, no. 6, 1991, pages 811-814, XP002082330 ISSN: 0040-4039 example 3 ----	1,2,9
X	EP 0 581 095 A (BASF AG) 2 February 1994 (1994-02-02) formula (I), examples 22-25 ----	1,2,6-9
X	US 4 607 035 A (BETTARINI FRANCO ET AL) 19 August 1986 (1986-08-19) formula (I); examples 1,11,14,28,32-34,37-39 ----	1,2,6-9
X	US 4 971 982 A (ATTWOOD MICHAEL R ET AL) 20 November 1990 (1990-11-20) formula (I); col. 23, l. 29; col. 36, l. 26; col. 47, l. 65; col. 50, l. 18 ----	1,2,6-9
X	EP 0 385 680 A (ICI PHARMA ; ICI PLC (GB)) 5 September 1990 (1990-09-05) formula (I); examples 3,4,6 -----	1,2,6-9



## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08878

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0194339	A	13-12-2001	AU 6234401 A WO 0194339 A1	17-12-2001 13-12-2001
WO 0228182	A	11-04-2002	AU 1820002 A WO 0228182 A1	15-04-2002 11-04-2002
WO 0155066	A	02-08-2001	AU 2680701 A WO 0155066 A2 EP 1250314 A2	07-08-2001 02-08-2001 23-10-2002
JP 11147866	A	02-06-1999	NONE	
DE 4115465	A	12-11-1992	DE 4115465 A1 AT 157974 T AU 1659192 A CA 2102904 A1 WO 9220671 A1 DE 59208898 D1 DK 586405 T3 EP 0586405 A1 ES 2106866 T3 GR 3025290 T3 JP 6508109 T PL 297368 A1	12-11-1992 15-09-1997 30-12-1992 12-11-1992 26-11-1992 16-10-1997 01-12-1997 16-03-1994 16-11-1997 27-02-1998 14-09-1994 20-09-1993
EP 0581095	A	02-02-1994	AU 4212193 A CA 2100546 A1 EP 0581095 A2 HU 66105 A2 JP 6211748 A NZ 248227 A ZA 9305332 A	27-01-1994 25-01-1994 02-02-1994 28-09-1994 02-08-1994 26-09-1995 23-01-1995
US 4607035	A	19-08-1986	IT 1218326 B BE 897354 A1 CA 1181403 A1 CH 660179 A5 DE 3326180 A1 FR 2530627 A1 GB 2124227 A , B JP 3026178 B JP 59033239 A NL 8302621 A	12-04-1990 23-01-1984 22-01-1985 31-03-1987 26-01-1984 27-01-1984 15-02-1984 10-04-1991 23-02-1984 16-02-1984
US 4971982	A	20-11-1990	AT 91127 T AU 613646 B2 AU 1855688 A CA 1334094 A1 CN 1030582 A , B CZ 8804841 A3 DE 3882095 D1 DK 370988 A EP 0298452 A2 ES 2056859 T3 FI 883232 A , B, HU 50152 A2 HU 9500257 A3 IE 62258 B	15-07-1993 08-08-1991 12-01-1989 24-01-1995 25-01-1989 13-10-1999 05-08-1993 07-01-1989 11-01-1989 16-10-1994 07-01-1989 28-12-1989 28-09-1995 11-01-1995

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08878

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4971982	A	IL 86923 A	27-02-1994
		JP 1038087 A	08-02-1989
		JP 2683581 B2	03-12-1997
		KR 9612213 B1	16-09-1996
		MC 1953 A	30-06-1989
		MX 12168 A ,B	01-10-1993
		NO 883007 A ,B,	09-01-1989
		NZ 225163 A	26-04-1991
		PH 27153 A	02-04-1993
		PT 87913 A ,B	30-06-1989
		SK 484188 A3	06-05-1998
		SU 1757466 A3	23-08-1992
		US 5118694 A	02-06-1992
		YU 130288 A1	30-06-1990
		ZA 8804656 A	06-01-1989
EP 0385680	A	05-09-1990	AU 625304 B2
			09-07-1992
			AU 4976090 A
			06-09-1990
			CA 2009900 A1
			31-08-1990
			DE 69020008 D1
			20-07-1995
			DE 69020008 T2
			05-10-1995
			EP 0385680 A2
			05-09-1990
			IE 66512 B1
			10-01-1996
			NO 900919 A
			29-08-1990
			NZ 232517 A
			28-04-1992
			PT 93290 A
			31-08-1990
			US 5202326 A
			13-04-1993
			ZA 9001046 A
			31-10-1990